

Flow decomposition for heat equations with memory

Gengsheng Wang*

Yubiao Zhang[†]

Enrique Zuazua[‡]

Abstract

We build up a decomposition for the flow generated by the heat equation with a real analytic memory kernel. It consists of three components: The first one is of parabolic nature; the second one gathers the hyperbolic component of the dynamics, with null velocity of propagation; the last one exhibits a finite smoothing effect. This decomposition reveals the hybrid parabolic-hyperbolic nature of the flow and clearly illustrates the significant impact of the memory term on the parabolic behavior of the system in the absence of memory terms.

Keywords. Heat equations with memory, decomposition of the flow, hybrid parabolic-hyperbolic behavior

1 Introduction

1.1 Statement of the problem

In this paper, we will study the following heat equation with memory:

$$\begin{cases} \partial_t y(t, x) - \Delta y(t, x) + \int_0^t M(t-s)y(s, x)ds = 0, & (t, x) \in \mathbb{R}^+ \times \Omega, \\ y(t, x) = 0, & (t, x) \in \mathbb{R}^+ \times \partial\Omega, \\ y(0, x) = y_0(x), & x \in \Omega. \end{cases} \quad (1.1)$$

Here, $\mathbb{R}^+ := (0, +\infty)$, $\Omega \subset \mathbb{R}^n$ ($n \in \mathbb{N}^+ := \{1, 2, 3, \dots\}$) is a bounded domain with a C^2 -boundary $\partial\Omega$, y_0 is an initial datum and M is a memory-kernel over $\overline{\mathbb{R}^+} := [0, +\infty)$.

Although our analysis can be generalized to less regular memory kernels, for the sake of simplicity we assume that:

(\mathfrak{C}) the memory kernel M is a real analytic and nonzero function over $\overline{\mathbb{R}^+}$.

*Center for Applied Mathematics, Tianjin University, Tianjin, 300072, China. e-mail: wanggs62@yeah.net

[†]Center for Applied Mathematics, Tianjin University, Tianjin, 300072, China; e-mail: yubiao.zhang@yeah.net

[‡][1] Chair in Applied Analysis, Alexander von Humboldt-Professorship, Department of Data Science, Friedrich-Alexander-Universität Erlangen-Nürnberg, 91058 Erlangen, Germany (e-mail: enrique.zuazua@fau.de),

[2] Chair of Computational Mathematics, Fundación Deusto, Av. de las Universidades, 24, 48007 Bilbao, Basque Country, Spain,

[3] Departamento de Matemáticas, Universidad Autónoma de Madrid, 28049 Madrid, Spain

Mathematics Subject Classification (2010): 45K05 35K05 93C05

Equations with memory arise in the modeling of many physical phenomena such as viscoelasticity, heat conduction, etc. They can be traced back to the works of J. Maxwell [14], L. Boltzmann [2, 3] and V. Volterra [19, 20]. For instance, in the analysis of elastic materials, L. Boltzmann and V. Volterra represented the stress tensor in terms of the strain tensor as well as its history values. Equations involving memory terms have been widely studied: see for instance [1, 4, 5, 6, 7, 8, 9, 10, 11, 13, 17] and the references therein. In particular, in [11] the general memory effect in heat conduction processes was analyzed showing that temperature waves travelling in the direction of the heat-flux propagate faster than wave travelling in the opposite direction, while in [8] the asymptotic behavior of the systems of linear viscoelasticity at large time was analyzed, introducing a new auxiliary variable to deal with the history of the states.

By standard methods (see, for instance, [18, Theorem 1.2 in Section 6.1, p. 184]), it can be shown that the equation (1.1), with $y_0 \in L^2(\Omega)$, has a unique mild solution, denoted by $y(\cdot; y_0)$, in the space $C(\overline{\mathbb{R}^+}; L^2(\Omega))$. For each $t \geq 0$, we let the evolution of the system be denoted by:

$$\Phi(t)y_0 := y(t; y_0), \quad y_0 \in L^2(\Omega). \quad (1.2)$$

For each $t \geq 0$, the flow generated by the equation (1.1), $\Phi(t)$, belongs to $\mathcal{L}(L^2(\Omega))$. Here and in what follows, we denote by $\mathcal{L}(E, F)$ (where E and F are two Banach spaces) the space of all linear and bounded operators from E to F , and simply write $\mathcal{L}(E)$ for $\mathcal{L}(E, E)$.

We shall use the notation $\{e^{tA}\}_{t \geq 0}$ for the C_0 semigroup generated by the heat equation in the absence of memory term (i.e., when $M \equiv 0$), where

$$Af := \Delta f, \quad \text{with its domain } D(A) := H^2(\Omega) \cap H_0^1(\Omega). \quad (1.3)$$

Then $z(t; y_0) := e^{tA}y_0$, $t \geq 0$, solves (1.1) without memory, i.e., (1.1) when $M \equiv 0$.

This paper is devoted to analyzing the dynamics of the system with memory term and, in particular, to exhibiting the significant differences with the heat semigroup in the absence of memory.

1.2 Main results

The aim of this paper is to build up a decomposition of the flow $\Phi(t)$, revealing a hybrid parabolic-hyperbolic dynamics of the (1.1).

To state our main results, we first introduce several concepts, definitions and notations.

- Let $\eta_j > 0$ be the j^{th} eigenvalue of $-A$ and let e_j be the corresponding normalized eigenfunction in $L^2(\Omega)$. Let $\eta_1 < \eta_2 \leq \dots \leq \eta_j \dots \rightarrow +\infty$. Define, for each $s \in \mathbb{R}$, the real Hilbert space:

$$\mathcal{H}^s := \left\{ f = \sum_{j=1}^{\infty} a_j e_j : (a_j)_{j \geq 1} \subset \mathbb{R}, \sum_{j=1}^{\infty} |a_j|^2 \eta_j^s < +\infty \right\}, \quad (1.4)$$

equipped with the inner product:

$$\langle f_1, f_2 \rangle_{\mathcal{H}^s} := \sum_{j=1}^{\infty} a_{j,1} a_{j,2} \eta_j^s, \quad f_k = \sum_{j=1}^{\infty} a_{j,k} e_j \in \mathcal{H}^s \quad (k = 1, 2).$$

For all $t \geq 0$, $\Phi(t)$ belongs to $\mathcal{L}(\mathcal{H}^s)$ for any $s \in \mathbb{R}$ (see Proposition 7.1).

We now introduce the classes:

$$\mathcal{H}^{-\infty} := \bigcup_{s \in \mathbb{R}} \mathcal{H}^s \quad \text{and} \quad \mathcal{H}^{+\infty} := \bigcap_{s \in \mathbb{R}} \mathcal{H}^s. \quad (1.5)$$

- Recall that for each continuous function f over \mathbb{R}^+ , the operator $f(-A)$ can be defined by the spectral functional calculus (see [15, Section 3, Chapter V.III]).
- Let us also define the following functions, related to the memory kernel M , that will play important roles in the decomposition of the flow.

First, we introduce the *flow kernel*:

$$K_M(t, s) := \sum_{j=1}^{+\infty} \frac{(-s)^j}{j!} \underbrace{M * \cdots * M}_j(t - s), \quad (t, s) \in S_+ := \{(t, s) \in \mathbb{R}^2 : t \geq s\}. \quad (1.6)$$

Here and throughout the paper, $*$ denotes the usual convolution, i.e., when $g_1, g_2 \in L_{loc}^1(\mathbb{R}^+)$,

$$g_1 * g_2(t) := \int_0^t g_1(t-s) g_2(s) ds, \quad t \geq 0.$$

Notice that the above K_M is well-defined and it is real analytic over S_+ (see Proposition 2.3). The following holds (see Proposition 4.8)

$$\Phi(t) = e^{tA} + \int_0^t K_M(t, \tau) e^{\tau A} d\tau, \quad t \geq 0,$$

which yields a clear description of the gap between the heat equation and the memory one and justifies the terminology “flow kernel” employed.

Second, for each $N \in \mathbb{N}^+$, let

$$\mathcal{R}_N(t, \tau) := \int_0^t \tau e^{-\tau s} \partial_s^N K_M(t, s) ds, \quad t \geq 0, \quad \tau \geq 0. \quad (1.7)$$

Third, we define two sequences of functions $\{h_l\}_{l \in \mathbb{N}}$ and $\{p_l\}_{l \in \mathbb{N}}$ (that will play the role of coefficients in the expansions) in the following manner: for each $t \geq 0$,

$$\begin{cases} h_l(t) := (-1)^l \sum_{j=0}^l C_l^{l-j} \frac{d^{(l-j)}}{dt^{(l-j)}} \underbrace{M * \cdots * M}_j(t); \\ p_l(t) := -h_l(0) + (-1)^{l+1} \sum_{\substack{m, j \in \mathbb{N}^+, \\ 2j-l-1 \leq m \leq j}} \left(C_l^{l-j+m} \frac{d^{(l-j+m)}}{dt^{(l-j+m)}} \underbrace{M * \cdots * M}_j(0) \right) \frac{(-t)^m}{m!}. \end{cases} \quad (1.8)$$

Here, $C_\beta^m := \beta!/m!(\beta-m)!$ and $\underbrace{M * \cdots * M}_j := 0$ when $j = 0$.

- Let f be a distribution over a domain $\mathcal{D} \subset \mathbb{R}^k$ (with $k \in \mathbb{N}^+$). By the notation $f \in L_{loc}^2(q)$ (with $q \in \mathcal{D}$), we refer to the fact that $f|_{\mathcal{U}_0} \in L^2(\mathcal{U}_0)$ for an open non-empty subset \mathcal{U}_0 such that $q \in \mathcal{U}_0 \subset \mathcal{D}$.

The main results of this paper are as follows.

Theorem 1.1. *For each integer $N \geq 2$, the flow $\Phi(t)$ admits the following decomposition:*

$$\Phi(t) = \mathcal{P}_N(t) + \mathcal{W}_N(t) + \mathfrak{R}_N(t), \quad t \geq 0, \quad (1.9)$$

with

$$\begin{cases} \mathcal{P}_N(t) &:= e^{tA} + e^{tA} \sum_{l=0}^{N-1} p_l(t)(-A)^{-l-1}, \\ \mathcal{W}_N(t) &:= \sum_{l=0}^{N-1} h_l(t)(-A)^{-l-1}, \\ \mathfrak{R}_N(t) &:= \mathcal{R}_N(t, -A)(-A)^{-N-1}, \end{cases} \quad t \geq 0, \quad (1.10)$$

where $\{h_l\}_{l \in \mathbb{N}}$ and $\{p_l\}_{l \in \mathbb{N}}$ are given by (1.8) and \mathcal{R}_N is given by (1.7). Moreover, for each $t \geq 0$, neither $\{h_l(t)\}_{l \geq 1}$ nor $\{p_l(t)\}_{l \in \mathbb{N}}$ is the null sequence, i.e.,

$$\sum_{l \geq 1} |h_l(t)| > 0 \quad \text{and} \quad \sum_{l \geq 1} |p_l(t)| > 0. \quad (1.11)$$

Theorem 1.2. *With the notations in Theorem 1.1, the following conclusions are true for each integer $N \geq 2$:*

- (i) *The first component \mathcal{P}_N exhibits a heat-like behavior: for each $t > 0$, $\mathcal{P}_N(t)\mathcal{H}^{-\infty} \subset \mathcal{H}^{+\infty}$, where $\mathcal{H}^{+\infty}$ and $\mathcal{H}^{-\infty}$ are given by (1.5).*
- (ii) *The second component \mathcal{W}_N exhibits a wave-like behavior: when $y_0 \in \mathcal{H}^{-\infty}$, $x_0 \in \Omega$ and $t_0 > 0$,*

$$\mathcal{W}_N(\cdot)y_0 \notin L_{loc}^2(t_0, x_0) \iff \forall t > 0, \mathcal{W}_N(\cdot)y_0 \notin L_{loc}^2(t, x_0). \quad (1.12)$$

In other words, the singularities of the solutions propagate in the time-like direction with null velocity of propagation in the space-like direction.

- (iii) *The last component \mathfrak{R}_N exhibits a time-uniform smoothing effect with a gain of $2N + 2$ space derivatives: for each $y_0 \in \mathcal{H}^s$ with $s \in \mathbb{R}$, $\mathfrak{R}_N(\cdot)y_0 \in C([0, +\infty); \mathcal{H}^{s+2N+2})$, while $A^{-j}e^{\cdot A}y_0, A^{-j}y_0 \in C([0, +\infty); \mathcal{H}^{s+2j})$ for any $0 \leq j \leq N$. And for each $s \in \mathbb{R}$, the term \mathcal{R}_N (in \mathfrak{R}_N) belongs to $C(\mathbb{R}^+; \mathcal{L}(\mathcal{H}^s))$ and fulfills the estimate:*

$$\|\mathcal{R}_N(t, -A)\|_{\mathcal{L}(\mathcal{H}^s)} \leq e^t \left\{ \exp \left[N(1+t) \left(\sum_{j=0}^N \max_{0 \leq \tau \leq t} \left| \frac{d^j}{d\tau^j} M(\tau) \right| \right) \right] - 1 \right\}, \quad t \geq 0. \quad (1.13)$$

(iv) For any $y_0 \in \mathcal{H}^{-\infty}$, $x_0 \in \Omega$ and $t > 0$,

$$\Phi(\cdot)y_0 \notin L_{loc}^2(t, x_0) \Leftrightarrow \mathcal{W}_N(\cdot)y_0 \notin L_{loc}^2(t, x_0) \Leftrightarrow A^{-2}y_0 \notin L_{loc}^2(x_0). \quad (1.14)$$

Theorem 1.3. *With the notations in Theorem 1.1, the following conclusions hold for each integer $N \geq 2$: First, at the initial time,*

$$\lim_{j \rightarrow +\infty} \|\Phi(0)e_j\|_{L^2(\Omega)} = \lim_{j \rightarrow +\infty} \|\mathcal{P}_N(0)e_j\|_{L^2(\Omega)} = 1, \quad \lim_{j \rightarrow +\infty} \|\mathcal{W}_N(0)e_j\|_{L^2(\Omega)} = 0, \quad \mathfrak{R}_N(0) = 0. \quad (1.15)$$

Second, at each time $t > 0$,

$$\begin{cases} \lim_{j \rightarrow +\infty} \|\Phi(t)e_j\|_{\mathcal{H}^4} = \lim_{j \rightarrow +\infty} \|\mathcal{W}_N(t)e_j\|_{\mathcal{H}^4} = |M(t)|, \\ \lim_{j \rightarrow +\infty} \|\Phi(t)e_j\|_{\mathcal{H}^s} = \lim_{j \rightarrow +\infty} \|\mathcal{W}_N(t)e_j\|_{\mathcal{H}^s} = 0 \text{ for any } s < 4, \\ \lim_{j \rightarrow +\infty} \|\mathcal{P}_N(t)e_j\|_{\mathcal{H}^s} = 0 \text{ for any } s \in \mathbb{R}, \\ \lim_{j \rightarrow +\infty} \|\mathfrak{R}_N(t)e_j\|_{\mathcal{H}^s} = 0 \text{ for any } s < 2N + 2. \end{cases} \quad (1.16)$$

Theorem 1.4. *Given $s \in \mathbb{R}$, the following conclusions are true:*

(i) *There is $C_0 > 0$ (independent of s) so that for any $\alpha \in [0, 4]$ and $t > 0$, $\Phi(t)$ belongs to $\mathcal{L}(\mathcal{H}^s, \mathcal{H}^{s+\alpha})$ and satisfies*

$$\|\Phi(t)\|_{\mathcal{L}(\mathcal{H}^s, \mathcal{H}^{s+\alpha})} \leq C_0 t^{-\frac{\alpha}{2}} \exp \left[2(1+t)(1 + \|M\|_{C^2([0,t])}) \right]. \quad (1.17)$$

(ii) *If there is $\alpha_0 \geq 0$ so that*

$$\Phi(t) \in \mathcal{L}(\mathcal{H}^s, \mathcal{H}^{s+\alpha_0}) \text{ as } t > 0 \text{ in a neighborhood of } 0, \quad (1.18)$$

then $\alpha_0 \leq 4$ and $\liminf_{t \rightarrow 0^+} t^{\frac{\alpha_0}{2}} \|\Phi(t)\|_{\mathcal{L}(\mathcal{H}^s, \mathcal{H}^{s+\alpha_0})} > 0$.

(iii) *For any $y_0 \in \mathcal{H}^s$, $\Phi(\cdot)y_0 \in C(\mathbb{R}^+; \mathcal{H}^{s+4})$. Moreover, the index 4 is optimal in the sense that if $\alpha > 4$, then $\Phi(\cdot)\hat{y}_0 \notin C(\mathbb{R}^+; \mathcal{H}^{s+\alpha})$ for some $\hat{y}_0 \in \mathcal{H}^s$.*

Remark 1.5. Several comments are in order:

(a1) Theorem 1.1 gives the decomposition (1.9) of the flow and besides shows the non-triviality of \mathcal{P}_N and \mathcal{W}_N : for each $t \geq 0$, there is $N_0(t) \in \mathbb{N}^+$ so that $\mathcal{P}_N(t) \neq 0$ and $\mathcal{W}_N(t) \neq 0$ when $N \geq N_0(t)$. Theorem 1.2 explains the functionality of each term in the decomposition (1.9). Theorems 1.3-1.4 are the consequences of the decomposition (1.9). The three terms of the decomposition \mathcal{P}_N , \mathcal{W}_N and \mathfrak{R}_N are referred to as the heat-like component, the wave-like component and the remainder respectively. The first two components in (1.9) are the main ones of the decomposition.

- (a2) The proof of Theorem 1.1 uses a Fourier expansion on the basis of eigenfunctions of A that reduces the problem to consider an ODE with memory depending on the dual parameter $\eta > 0$:

$$w'(t) + \eta w(t) + \int_0^t M(t-s)w(s)ds = 0, \quad t > 0; \quad w(0) = 1. \quad (1.19)$$

The dynamics of this memory-ODE can be decomposed into three terms leading to the three components in the decomposition (1.9). The asymptotics of this decomposition for large η yields the main properties of the decomposition (1.9) of the memory-heat equation. A careful analysis of the flow kernel K_M plays a key role in this analysis.

- (a3) In Theorem 1.2, the infinite order regularizing effect of the heat-like behavior of \mathcal{P}_N , stated in (i), is the analog of the smoothing effect of the semigroup $\{e^{tA}\}_{t \geq 0}$ generated by the heat equation without memory terms; The wave-like component \mathcal{W}_N exhibits the propagation of singularities along the time-direction, as stated in (ii); The smoothing effect of the remainder \mathfrak{R}_N , stated in (iii), ensures the gain of $2N + 2$ space-derivatives at nonnegative time but differs from the infinite order smoothing effect of the heat semigroup e^{tA} at positive time (see Remark 4.4 for more discussions). The conclusion (iv) says, in plain language, that when $t > 0$, the singularity of $\mathcal{W}_N(t)$ determines the singularity of $\Phi(t)$, more precisely, the singularity of the practical leading term in $\mathcal{W}_N(t)$ determines the singularity of $\Phi(t)$. Here, we notice that it follows by (1.8) that $h_0(t) \equiv 0$ and $h_1(t) = -M(t)$, thus the practical leading term in $\mathcal{W}_N(t)$ is $-M(t)A^2$ where the coefficient $M(t)$ is not zero except for finitely many t by the assumption (C).

From these, we conclude that the decomposition (1.9) in Theorem 1.1 reveals the hybrid parabolic-hyperbolic behavior of the flow $\Phi(t)$.

- (a4) Theorem 1.3 shows how the energy of solutions taking eigenfunctions of the operator A as initial data is distributed over each component of (1.9) at time $t = 0$ and time $t > 0$ respectively. The conclusion (1.15) says that when $t = 0$, the energy of $\Phi(0)e_j$ (which is exactly e_j) is almost concentrated in the heat-like component for large j , while (1.16) can be explained as: when $t > 0$, the energy of $\Phi(t)e_j$ almost focuses on the wave-like component for large j . The first line in (1.16) is from the term $-M(t)A^{-2}$ (in \mathcal{W}_N) and the order 4 in \mathcal{H}^4 is exactly from the order -2 in A^{-2} . The last line in (1.16) is from (iii) in Theorem 1.2.
- (a5) Theorem 1.4 exhibits the finite order smoothing effect of the flow. By it, we can see, from the point of view of the smoothing effect, both big differences and some similarities between the flow $\Phi(t)$ and the semigroup e^{tA} :

First, on one hand, for any $y_0 \in \mathcal{H}^s$ (with $s \in \mathbb{R}$), $e^{tA}y_0 \in C(\mathbb{R}^+; \mathcal{H}^k)$ ($\forall k \in \mathbb{N}$), while $\Phi(\cdot)y_0 \in C(\mathbb{R}^+; \mathcal{H}^{s+4})$ and moreover the index 4 is optimal, on the other hand, the

smoothing effect of the flow $\Phi(t)$ at points in the set $\{t > 0 : M(t) = 0\}$ is better than that at points in the set $\{t > 0 : M(t) \neq 0\}$ (see Remark 4.5 for more details); Second, when t is large, both the semigroup e^{tA} and the flow $\Phi(t)$ are bounded from the above by exponential functions of t , while when t is small, they are bounded from the above by the function t^{-2} . Moreover, the flow $\Phi(t)$ is also bounded from the below by t^{-2} when $t \rightarrow 0^+$. Here, by “ $e^{tA}/\Phi(t)$ is bounded by”, we mean “the $\mathcal{L}(\mathcal{H}^s, \mathcal{H}^{s+4})$ -norm of $e^{tA}/\Phi(t)$ is bounded by”.

- (a6) It follows from Theorem 1.1 that for each $t \geq 0$, the wave-like component $\mathcal{W}_N(t)$, as well as the gap between the heat-like component $\mathcal{P}_N(t)$ and the heat semigroup e^{tA} , are non-trivial, when N is large enough. Thus, we may expect that as N increases the heat-like and the wave-like components include an increasing number of terms, just like in the Taylor expansion, and the decomposition becomes sharper. (See the example in Section 5.)
- (a7) We require $N \geq 2$ in Theorem 1.1 since $\mathcal{W}_1(\cdot) \equiv 0$ (when $N = 1$).
- (a8) Notice that both $\mathcal{P}_N(0)$ and $\mathcal{W}_N(0)$ are not projection operators in general and that

$$y_0 = \mathcal{P}_N(0)y_0 + \mathcal{W}_N(0)y_0 \text{ for each } y_0 \in L^2(\Omega) \text{ and } \mathfrak{R}_N(0) = 0.$$

This is further discussed in Proposition 4.7.

- (a9) Whether a complete meaningful decomposition can be achieved without the reminder term is an interesting open problem (see Section 6).
- (a10) All the results of Theorems 1.1-1.2, except (1.11), (1.12) and (1.14), hold under the weaker assumption $M \in C^\infty(\overline{\mathbb{R}^+})$. In Theorem 4.10 below we analyse the case of kernels M in $C^{N_0}(\overline{\mathbb{R}^+})$, with $N_0 \geq 2$. The assumption (C) ensures (1.11), (1.12) and (1.14). Whether the same holds under weaker conditions on the kernel is an open problem, see Section 6.
- (a11) The decomposition (1.9) has applications in control theory. It allows, in particular, to compare the reachable sets for the controlled heat equations with and without memory term. We refer to [21] for a complete analysis of this issue.
- (a12) There is a large body of literature on the large time dynamics of memory like problems (see, for instance, [8, 9]) which is surely an important direction. Unfortunately, we are not able to use our decomposition to get such results.

1.3 Plan of the paper

The rest of the paper is organized as follows. In Section 2 we analyze the flow kernel K_M . In Section 3 we present a decomposition for solutions to the ODE (1.19). Section 4 contains the proofs of Theorems 1.1-1.4, and provides some other properties of the flow. In Section 5 we discuss, as an example, the case of the kernel $M(t) = \alpha e^{\lambda t}$. Section 6 lists several open problems. Section 7 contains an appendix.

2 Properties of the flow kernel

This section presents some properties of the flow kernel K_M in (1.6), which will be used later.

In what follows, the space $C^k([a, b])$ (with $k \in \mathbb{N}^+$ and $a < b$) is endowed with the norm:

$$\|f\|_{C^k([a, b])} := \sum_{l=0}^k \left\| \frac{d^l f}{dx^l} \right\|_{C([a, b])}, \quad f \in C^k([a, b]).$$

The following result provides basic estimates on iterated convolutions that will be used in the proof of Proposition 2.2. Its proof is put in the appendix.

Lemma 2.1. *Let $j, m \in \mathbb{N}^+$. Then for each sequence $\{M_l\}_{l=1}^j \subset C^m(\overline{\mathbb{R}^+})$, $M_1 * \dots * M_j$ is in the space $C^{m+j-1}(\overline{\mathbb{R}^+})$ and satisfies that for each $k \in \{0, 1, \dots, m+j-1\}$,*

$$\left| \frac{d^k}{dt^k} M_1 * \dots * M_j(t) \right| \leq \left(\sum_{l=\max\{0, j-1-k\}}^{j-1} \frac{t^l}{l!} \right) \prod_{l=1}^j \|M_l\|_{C^p([0, t])}, \quad t > 0, \quad (2.1)$$

where $p = \chi_{\mathbb{N}}(k-1)[(k-j)\chi_{\mathbb{N}}(k-j)+1]$ and $\chi_{\mathbb{N}}$ is the characteristic function of the set \mathbb{N} .

The following Proposition 2.2 provides estimates on the derivatives of the flow kernel K_M that will serve for the proof of Theorem 3.1, which is one of the tools in the proof of the decomposition in Theorem 1.1.

Proposition 2.2. *The flow kernel $K_M \in C^\infty(S_+)$ (where S_+ is given in (1.6)) satisfies that for each $\alpha, \beta \in \mathbb{N}$,*

$$|\partial_t^\alpha \partial_s^\beta K_M(t, s)| \leq e^{t-s} \left[\exp \left(\beta(1+|s|) \|M\|_{C^{\alpha+\beta}([0, t-s])} \right) - 1 \right], \quad t > s. \quad (2.2)$$

Proof. First of all, it follows from (1.6) that

$$K_M(t, s) = \sum_{j=1}^{+\infty} \mathcal{M}_j(t, s), \quad t \geq s, \quad (2.3)$$

where

$$\mathcal{M}_j(t, s) := \frac{(-s)^j}{j!} \underbrace{M * \dots * M}_j(t-s), \quad t \geq s. \quad (2.4)$$

Next, we prove $K_M \in C^\infty(S_+)$ showing the convergence in $C^\infty(S_+)$ of the series on the right-hand side of (2.3).

To this end, we will estimate \mathcal{M}_j with $j \in \mathbb{N}^+$: By the assumption (\mathfrak{C}) and Lemma 2.1, we see that $\underbrace{M * \dots * M}_j$ belongs to $C^\infty(\overline{\mathbb{R}^+})$ and satisfies, for each $k \in \mathbb{N}$,

$$\left| \frac{d^k}{d\tau^k} \underbrace{M * \dots * M}_j(\tau) \right| \leq e^\tau \|M\|_{C^k([0, \tau])}^j, \quad \tau > 0. \quad (2.5)$$

From (2.4) it follows that $\mathcal{M}_j \in C^\infty(S_+)$.

By direct computations, for $\alpha, \beta \in \mathbb{N}$, we have $t \geq s$,

$$\begin{aligned}
\partial_t^\alpha \partial_s^\beta \mathcal{M}_j(t, s) &= \partial_s^\beta \partial_t^\alpha \mathcal{M}_j(t, s) \\
&= \frac{d^\beta}{ds^\beta} \left[\frac{(-s)^j}{j!} \left(\frac{d^\alpha}{d\tau^\alpha} \underbrace{M * \cdots * M}_j(\tau) \right) \Big|_{\tau=t-s} \right] \\
&= \sum_{m=0}^{\beta} C_\beta^m \frac{d^m}{d\tau^m} \left(\frac{(-s)^j}{j!} \right) (-1)^{\beta-m} \left(\frac{d^{\beta-m}}{d\tau^{\beta-m}} \frac{d^\alpha}{d\tau^\alpha} \underbrace{M * \cdots * M}_j(\tau) \right) \Big|_{\tau=t-s} \\
&= \frac{(-1)^\beta}{j!} \sum_{m=0}^{\min\{\beta, j\}} C_\beta^m (C_j^m m!) (-s)^{j-m} \left(\frac{d^{\alpha+\beta-m}}{d\tau^{\alpha+\beta-m}} \underbrace{M * \cdots * M}_j(\tau) \right) \Big|_{\tau=t-s}. \quad (2.6)
\end{aligned}$$

Here and in what follows, we use the conventional notation $0^0 = 1$. By (2.6) and (2.5), one has that, when $t > s$,

$$\begin{aligned}
\left| \partial_t^\alpha \partial_s^\beta \mathcal{M}_j(t, s) \right| &\leq \frac{1}{j!} \sum_{m=0}^j (C_\beta^m m!) C_j^m |s|^{j-m} \left(e^{t-s} \|M\|_{C^{\alpha+\beta}([0, t-s])}^j \right) \\
&\leq \frac{1}{j!} \beta^j \left(\sum_{m=0}^j C_j^m |s|^{j-m} \right) e^{t-s} \|M\|_{C^{\alpha+\beta}([0, t-s])}^j \\
&= \frac{e^{t-s}}{j!} \left(\beta(1 + |s|) \|M\|_{C^{\alpha+\beta}([0, t-s])} \right)^j. \quad (2.7)
\end{aligned}$$

Now, by (2.7) it follows that the series in (2.3) converges in $C^\infty(S_+)$.

Finally, by (2.3) and (2.7), after direct computations, we see that when $t > s$,

$$\left| \partial_t^\alpha \partial_s^\beta K_M(t, s) \right| \leq \sum_{j=1}^{\infty} \left| \partial_t^\alpha \partial_s^\beta \mathcal{M}_j(t, s) \right| \leq e^{t-s} \left[\exp \left(\beta(1 + |s|) \|M\|_{C^{\alpha+\beta}([0, t-s])} \right) - 1 \right].$$

This gives the desired estimate (2.2) and ends the proof of Proposition 2.2. \square

The next Proposition 2.3 concerns the analyticity of the flow kernel K_M . It will be used in the proofs of Proposition 2.4 and Proposition 4.9.

Proposition 2.3. *The flow kernel K_M is real analytic on S_+ (where S_+ is given in (1.6)).*

Proof. It suffices to prove that K_M is real analytic over S_T for each $T > 0$, where

$$S_T := \{(t, s) \in \mathbb{R}^2 : 0 \leq t - s \leq T\}.$$

To this end, we fix an arbitrary $T > 0$. Due to the analyticity of M over $\overline{\mathbb{R}^+}$, there is a domain $\widetilde{\mathcal{O}}_T$ of the complex plane \mathbb{C} , with $[0, T] \subset \widetilde{\mathcal{O}}_T \subset \mathbb{C}$, so that M has a unique analytic extension \widetilde{M} to $\widetilde{\mathcal{O}}_T$. Moreover, we can take a bounded and convex subdomain \mathcal{O}_T so that $[0, T] \subset \mathcal{O}_T \subset \subset \widetilde{\mathcal{O}}_T$. The convolution $*$ can then be extended to $\tilde{*}$ in the following manner for $f, g \in C(\mathcal{O}_T; \mathbb{C})$,

$$f \tilde{*} g(z) := \int_0^1 f((1-s)z) g(sz) z ds, \quad z \in \mathcal{O}_T.$$

We now claim the following two properties:

(P1) For each $j \in \mathbb{N}^+$, $\underbrace{\widetilde{M} \tilde{*} \cdots \tilde{*} \widetilde{M}}_j$ is an analytic extension of $\underbrace{M * \cdots * M}_j$ over \mathcal{O}_T ;

(P2) There is $C > 0$ so that

$$\sup_{z \in \mathcal{O}_T} |\underbrace{\widetilde{M} \tilde{*} \cdots \tilde{*} \widetilde{M}}_j(z)| \leq C^j \text{ for all } j \in \mathbb{N}^+.$$

Indeed, one has

$$f \tilde{*} g|_{[0,T]} = f|_{[0,T]} * g|_{[0,T]}, \text{ when } f, g \in C(\mathcal{O}_T; \mathbb{C}),$$

Here, $f \tilde{*} g|_{[0,T]}$, $f|_{[0,T]}$ and $g|_{[0,T]}$ are respectively the restrictions of $f \tilde{*} g$, f and g over $[0, T]$. Then, property (P1) follows from the analyticity of M at once, while the property (P2) can be proved by direct computations.

Define the following subset of \mathbb{C}^2 :

$$\mathcal{D}_T := \left\{ (t, s) \in \mathbb{C}^2 : t - s \in \mathcal{O}_T \right\}.$$

It is clear that $S_T \subset \mathcal{D}_T$. According to (P2) above, the following series uniformly converges over each compact subset of \mathcal{D}_T :

$$\sum_{j \geq 1} \frac{(-s)^j}{j!} \underbrace{\widetilde{M} \tilde{*} \cdots \tilde{*} \widetilde{M}}_j(t - s), \quad (t, s) \in \mathcal{D}_T.$$

Meanwhile, by (P1), we find that each term in the above series is analytic over \mathcal{D}_T . Hence, the sum of this series is analytic over \mathcal{D}_T . From this and (1.6), we see that $K_M|_{S_T}$ (the restriction of K_M over S_T) can be analytically extended to \mathcal{D}_T . Therefore, it is real analytic over S_T . This ends the proof of Proposition 2.3. \square

The next Proposition 2.4 can be interpreted as a strong unique continuation property or non-degeneracy of the kernel K_M , that will be used in the proof of Theorem 3.1.

Proposition 2.4. *For each $(t_0, s_0, v_1, v_2) \in (S_+ \times \mathbb{S}^1) \setminus I$, where*

$$I := \left\{ (t, t, \tau, \tau) : t \in \mathbb{R}, \tau = \pm 1/\sqrt{2} \right\} \cup \left\{ (t, 0, \tau, 0) : t \geq 0, \tau = \pm 1 \right\},$$

with $\partial_{\vec{v}} := v_1 \partial_t + v_2 \partial_s$ where $\vec{v} := (v_1, v_2)$, it holds that

$$\partial_{\vec{v}}^l K_M(t_0, s_0) \neq 0 \text{ for some } l \in \mathbb{N}. \quad (2.8)$$

Remark 2.5. *Obviously, real analytic function on \mathbb{R}^2 do not necessarily fulfill the non-degeneracy condition above. Indeed, polynomials, for instance, can vanish along lines in \mathbb{R}^2 .*

Proof of Proposition 2.4. It suffices to prove (2.8) in the case that $\vec{v} = (0, 1)$, as other cases can be proved in a very similar way. By contradiction, suppose that (2.8) with $\vec{v} = (0, 1)$ fails, i.e.,

$$\partial_{(0,1)}^l K_M(t_0, s_0) = 0 \text{ for all } l \in \mathbb{N}. \quad (2.9)$$

Define

$$f(\lambda) := K_M(t_0, t_0 - \lambda), \quad \lambda \geq 0. \quad (2.10)$$

Two facts on f are given as follows: First, by the real analyticity of M over $\overline{\mathbb{R}^+}$, we see from Proposition 2.3 that K_M is real analytic over S_+ . This, along with (2.10), yields that f is real analytic over $\overline{\mathbb{R}^+}$. Second, by (2.9) and (2.10), we find that f vanishes of infinite order at $\lambda = t_0 - s_0$. From these two facts, we see that $f \equiv 0$ over $\overline{\mathbb{R}^+}$, which, along with (2.10), yields

$$K_M(t_0, t_0 - \lambda) = 0, \quad \lambda \geq 0.$$

The above, together with (1.6), shows

$$0 = \sum_{j=1}^{+\infty} \frac{(\lambda - t_0)^j}{j!} \underbrace{M * \cdots * M}_j(\lambda), \quad \lambda \geq 0,$$

which leads to

$$0 = M(\lambda) + \sum_{j=2}^{+\infty} \frac{(\lambda - t_0)^{j-1}}{j!} \underbrace{M * \cdots * M}_j(\lambda), \quad \lambda \geq 0. \quad (2.11)$$

Next, we arbitrarily fix $T > 0$. For each $j \in \mathbb{N}^+ \setminus \{1\}$, we define an operator \mathcal{K}_j on $C([0, T])$ in the following manner: given $g \in C([0, T])$, set

$$\mathcal{K}_j(g)(\lambda) := \frac{(\lambda - t_0)^{j-1}}{j!} \underbrace{M * \cdots * M}_{j-1} * g(\lambda), \quad 0 \leq \lambda \leq T, \quad (2.12)$$

which is well-defined, linear and bounded. Let

$$\mathcal{Q} := \sum_{j=2}^{+\infty} \mathcal{K}_j.$$

One can directly check that $\mathcal{Q} \in \mathcal{L}(C([0, T]))$, the Banach space of all linear and bounded operators on $C([0, T])$. Thus, we deduce from (2.11) that

$$(Id + \mathcal{Q})(M|_{[0, T]}) = \left(Id + \sum_{j=2}^{+\infty} \mathcal{K}_j \right) (M|_{[0, T]}) = 0, \quad (2.13)$$

where Id is the identity operator on $C([0, T])$.

We now claim that $(Id + \mathcal{Q})^{-1}$ exists in $\mathcal{L}(C([0, T]))$. When this is done, we can use (2.13), (1.6) and the arbitrariness of T to see that $M \equiv 0$, which contradicts the assumption (\mathfrak{C}) . Consequently, (2.8) is true.

The remainder is to show the above claim. To this end, we arbitrarily fix $k \in \mathbb{N}^+$ and then estimate \mathcal{Q}^k in the following manner: Set

$$T_{t_0} := \sup_{0 \leq \lambda \leq T} |\lambda - t_0| = \max \{ |t_0|, |T - t_0| \}.$$

Then from (2.12), one can directly check that when $j_1, \dots, j_k \geq 2$,

$$\begin{aligned} \|\mathcal{K}_{j_1} \cdots \mathcal{K}_{j_k}\|_{\mathcal{L}(C([0,T]))} &\leq \frac{1}{j_1!} \cdots \frac{1}{j_k!} T_{t_0}^{j_1+\dots+j_k-k} \underbrace{\| |M| * \cdots * |M| \|}_{j_1+\dots+j_k-k} \|_{L^1([0,T])} \\ &\leq \frac{1}{j_1!} \cdots \frac{1}{j_k!} T_{t_0}^{j_1+\dots+j_k-k} \frac{T_{t_0}^{j_1+\dots+j_k-k}}{(j_1+\dots+j_k-k)!} \|M\|_{C([0,T])}^{j_1+\dots+j_k-k} \\ &\leq \frac{1}{k!} \frac{1}{j_1!} \cdots \frac{1}{j_k!} \left(T_{t_0} T \|M\|_{C([0,T])} \right)^{j_1+\dots+j_k-k}. \end{aligned}$$

This, along with the definition of \mathcal{Q} , yields

$$\begin{aligned} \|\mathcal{Q}^k\|_{\mathcal{L}(C([0,T]))} &= \left\| \sum_{j_1, \dots, j_k \geq 2} \mathcal{K}_{j_1} \cdots \mathcal{K}_{j_k} \right\|_{\mathcal{L}(C([0,T]))} \\ &\leq \frac{1}{k!} \prod_{m=1}^k \left(\sum_{j_m \geq 2} \frac{1}{j_m!} \left(T_{t_0} T \|M\|_{C([0,T])} \right)^{j_m-1} \right) \leq \frac{1}{k!} \exp \left(k T_{t_0} T \|M\|_{C([0,T])} \right). \end{aligned}$$

So $\sum_{k=0}^{+\infty} (-\mathcal{Q})^k$ converges in $\mathcal{L}(C([0,T]))$. Then we have

$$Id = (Id + \mathcal{Q}) \sum_{k=0}^{+\infty} (-\mathcal{Q})^k.$$

Therefore, $(Id + \mathcal{Q})^{-1}$ exists in $\mathcal{L}(C([0,T]))$. This completes the proof of Proposition 2.4. \square

The following Proposition 2.6 presents a weighted estimate of the flow kernel K_M . It will be used in the proof of Proposition 4.8 that provides an explicit expression of the gap between the heat evolution with and without memory.

Proposition 2.6. *For each $\lambda \in \mathbb{R}$,*

$$\int_0^t e^{-\lambda(t-s)} |K_M(t, s)| ds \leq \exp \left(t \int_0^t e^{-\lambda\tau} |M(\tau)| d\tau \right) - 1, \quad t \geq 0. \quad (2.14)$$

Proof. Arbitrarily fix $\lambda \in \mathbb{R}$. Define the following weighted memory kernel:

$$M_\lambda(t) := e^{-\lambda t} M(t), \quad t \geq 0. \quad (2.15)$$

We claim

$$e^{-\lambda(t-s)} K_M(t, s) = \sum_{j=1}^{+\infty} \frac{(-s)^j}{j!} \underbrace{M_\lambda * \cdots * M_\lambda}_j(t-s), \quad (t, s) \in S_+. \quad (2.16)$$

Indeed, (2.16) follows from (1.6) and the following identity:

$$\underbrace{M_\lambda * \cdots * M_\lambda}_j(\tau) = e^{-\lambda\tau} \underbrace{M * \cdots * M}_j(\tau), \quad \tau \geq 0, \quad j \in \mathbb{N}^+,$$

which can be verified directly.

Next, we arbitrarily fix $t > 0$. By the iterative use of the Young's inequality:

$$\|f * g\|_{L^1(0,t)} \leq \|f\|_{L^1(0,t)} \|g\|_{L^1(0,t)}, \text{ when } f, g \in L^1(0,t),$$

one has that

$$\underbrace{\|M_\lambda * \cdots * M_\lambda\|_{L^1(0,t)}}_j \leq \|M_\lambda\|_{L^1(0,t)}^j, \quad j \in \mathbb{N}^+.$$

This, along with (2.16), yields

$$\int_0^t |e^{-\lambda(t-s)} K_M(t,s)| ds \leq \sum_{j=1}^{+\infty} \frac{t^j}{j!} \underbrace{\|M_\lambda * \cdots * M_\lambda\|_{L^1(0,t)}}_j \leq \exp\left(t \|M_\lambda\|_{L^1(0,t)}\right) - 1.$$

Then (2.14) follows from (2.15). This concludes the proof of Proposition 2.6. \square

3 Parametrized ODEs with memory

This section studies the ODE (1.19), i.e.,

$$w'(t) + \eta w(t) + \int_0^t M(t-s)w(s)ds = 0, \quad t > 0; \quad w(0) = 1, \quad (3.1)$$

where $\eta > 0$ is a parameter. First of all, by a standard method in the ODE theory, one can easily check that the equation (3.1) has a unique solution, denoted by w_η , in the space $C^1(\overline{\mathbb{R}^+})$. The main result of this section is the next Theorem 3.1 which gives a decomposition in terms of η for the solution w_η . It lays a solid foundation for the proof of Theorem 1.1.

Theorem 3.1. *For each integer $N \geq 2$, the solution w_η (with $\eta > 0$) to the equation (3.1) satisfies*

$$w_\eta(t) = e^{-\eta t} \left(1 + \sum_{l=0}^{N-1} p_l(t) \eta^{-l-1} \right) + \sum_{l=0}^{N-1} h_l(t) \eta^{-l-1} + \mathcal{R}_N(t, \eta) \eta^{-N-1}, \quad t \geq 0, \quad (3.2)$$

where $\{h_l\}_{l \in \mathbb{N}}$ and $\{p_l\}_{l \in \mathbb{N}}$ are given by (1.8) and \mathcal{R}_N is given by (1.7). In addition, the following conclusions are true:

(i) The function \mathcal{R}_N is in the space $C(\overline{\mathbb{R}^+} \times \overline{\mathbb{R}^+}) \cap C(\mathbb{R}^+; C(\mathbb{R}^+))$ and fulfills the estimate:

$$\|\mathcal{R}_N(t, \cdot)\|_{C(\mathbb{R}^+)} \leq e^t \left\{ \exp \left[N(1+t) \left(\sum_{j=0}^N \max_{0 \leq s \leq t} \left| \frac{d^j}{ds^j} M(s) \right| \right) \right] - 1 \right\}, \quad t \geq 0. \quad (3.3)$$

(ii) For each $t \in \overline{\mathbb{R}^+}$, neither $\{h_l(t)\}_{l \geq 1}$ nor $\{p_l(t)\}_{l \in \mathbb{N}}$ is the null sequence.

Remark 3.2. (i) The decomposition (3.2) serves for that in (1.9). The components in the decompositions (3.2) and (1.9) correspond to each other.

(ii) The conclusion (ii) in Theorem 3.1 corresponds to the conclusion (v) in Theorem 1.1 discussed in Remark 1.5.

To prove Theorem 3.1, we need several lemmas. The first one refers to the representation of the solutions of (3.1) in terms of the flow kernel K_M (given by (1.6)).

Lemma 3.3. *The solution w_η (with $\eta \in \mathbb{R}$) to (3.1) satisfies*

$$w_\eta(t) = e^{-\eta t} + \int_0^t K_M(t, s) e^{-\eta s} ds, \quad t \geq 0. \quad (3.4)$$

Proof. Fix an arbitrary $\eta \in \mathbb{R}$. From (3.1), it follows that

$$w'_\eta(t) + \eta w_\eta(t) = -M * w_\eta(t), \quad t \geq 0; \quad w_\eta(0) = 1.$$

The above yields

$$w_\eta(t) = e^{-\eta t} - \int_0^t e^{-\eta(t-s)} (M * w_\eta)(s) ds = e^{-\eta t} - (e^{-\eta \cdot} * M * w_\eta)(t), \quad t \geq 0. \quad (3.5)$$

Then we arbitrarily fix $T > 0$ and define the following operator:

$$\mathcal{Q}_T(f) := e^{-\eta \cdot} * M * f \quad \text{for each } f \in C([0, T]). \quad (3.6)$$

One can easily check that \mathcal{Q}_T is a linear and bounded operator on $C([0, T])$. By (3.6), we see that for each $k \in \mathbb{N}^+$,

$$\mathcal{Q}_T^k(f) = \underbrace{(e^{-\eta \cdot} * M) * \cdots * (e^{-\eta \cdot} * M)}_k * f, \quad f \in C([0, T]). \quad (3.7)$$

From (3.7), one can directly check

$$\begin{aligned} \|\mathcal{Q}_T^k\|_{\mathcal{L}(C([0, T]))} &\leq \|\underbrace{(e^{-\eta \cdot} * M) * \cdots * (e^{-\eta \cdot} * M)}_k\|_{L^1(0, T)} \\ &\leq \|e^{-\eta \cdot} * M\|_{C([0, T])}^k \left(\int_0^T \int_0^{t_1} \cdots \int_0^{t_{k-1}} dt_k \cdots dt_1 \right) \\ &\leq \left(e^{T|\eta|} \|M\|_{L^1(0, T)} \right)^k \frac{T^k}{k!}. \end{aligned}$$

From this, we see that the series $\sum_{k=0}^\infty (-\mathcal{Q}_T)^k$ converges in $\mathcal{L}(C([0, T]))$ and that

$$(1 + \mathcal{Q}_T)^{-1} = \sum_{k=0}^\infty (-\mathcal{Q}_T)^k. \quad (3.8)$$

Meanwhile, it follows from (3.5) and (3.6) that

$$w_\eta|_{[0, T]} = -\mathcal{Q}_T(w_\eta|_{[0, T]}) + e^{-\eta \cdot}|_{[0, T]}. \quad (3.9)$$

Here, $w_\eta|_{[0, T]}$ and $e^{-\eta \cdot}|_{[0, T]}$ denote respectively the restrictions of w_η and $e^{-\eta \cdot}$ over $[0, T]$.

Now, by (3.9) and (3.8), we find

$$w_\eta(t) = \left(\left(\sum_{j=0}^\infty (-\mathcal{Q}_T)^j \right) (e^{-\eta \cdot}|_{[0, T]}) \right)(t), \quad t \in [0, T],$$

from which and (3.7), it follows that

$$\begin{aligned} w_\eta(t) &= e^{-\eta t} + \sum_{j=1}^{\infty} (-1)^j \underbrace{(e^{-\eta \cdot} * M) * \cdots * (e^{-\eta \cdot} * M)}_j * e^{-\eta \cdot}(t) \\ &= e^{-\eta t} + \sum_{j=1}^{\infty} (-1)^j \underbrace{e^{-\eta \cdot} * \cdots * e^{-\eta \cdot}}_{j+1} * \underbrace{M * \cdots * M}_j(t), \quad t \in [0, T]. \end{aligned}$$

This, together with the following equality:

$$\underbrace{e^{-\eta \cdot} * \cdots * e^{-\eta \cdot}}_{j+1}(t) = e^{-\eta t} t^j / j!, \quad t \geq 0, \quad j \in \mathbb{N}^+,$$

shows

$$w_\eta(t) = e^{-\eta t} + \sum_{j=1}^{\infty} \int_0^t e^{-\eta s} \frac{(-s)^j}{j!} \underbrace{M * \cdots * M}_j(t-s) ds, \quad t \in [0, T].$$

Since $T > 0$ was arbitrarily taken, the above, along with (1.6), leads to (3.4). This ends the proof of Lemma 3.3. \square

The next lemma will be used to get an asymptotic expansion of the second term on the right-hand side of (3.4).

Lemma 3.4. *Given an integer $N \geq 2$ and a number $\eta > 0$, the following equality is true:*

$$\begin{aligned} \int_0^t e^{-\eta s} K_M(t, s) ds &= e^{-\eta t} \sum_{l=0}^{N-1} \left(-\partial_s^l K_M(t, s) \Big|_{s=t} \right) \eta^{-l-1} + \sum_{l=0}^{N-1} \left(\partial_s^l K_M(t, s) \Big|_{s=0} \right) \eta^{-l-1} \\ &\quad + \eta^{-N} \int_0^t e^{-\eta s} \partial_s^N K_M(t, s) ds, \quad t \geq 0. \end{aligned} \quad (3.10)$$

Proof. Fix $N \geq 2$, $\eta > 0$ and $t > 0$ arbitrarily. Given $g \in C([0, t])$, we define

$$\mathcal{F}_\eta(g) := \int_0^t e^{-\eta s} g(s) ds. \quad (3.11)$$

We first claim

$$\mathcal{F}_\eta(g) = \eta^{-N} \mathcal{F}_\eta(g^{(N)}) + \sum_{l=0}^{N-1} \eta^{-l-1} \left(g^{(l)}(0) - e^{-\eta t} g^{(l)}(t) \right), \quad \text{when } g \in C^N([0, t]). \quad (3.12)$$

Given $g \in C^N([0, t])$, from (3.11) and using the integration by parts, we find

$$\begin{aligned} \mathcal{F}_\eta(g) &= (-\eta)^{-1} \int_0^t \frac{d}{ds} e^{-\eta s} g(s) ds \\ &= (-\eta)^{-1} \left(e^{-\eta s} g(s) \right) \Big|_{s=0}^t + \frac{1}{\eta} \int_0^t e^{-\eta s} g'(s) ds \\ &= \eta^{-1} \left(g(0) - e^{-\eta t} g(t) \right) + \eta^{-1} \mathcal{F}_\eta(g'). \end{aligned}$$

Now, the iterative use of the above equality (to the derivatives of g) gives (3.12).

Next, it follows by Proposition 2.2 that $K_M(t, \cdot) \in C^\infty([0, t])$. Thus we can apply (3.12) (where $g(s) = K_M(t, s)$, $0 \leq s \leq t$) to get

$$\begin{aligned} \int_0^t K_M(t, s) e^{-\eta s} ds &= \eta^{-N} \int_0^t e^{-\eta s} \partial_s^N K_M(t, s) ds \\ &\quad + \sum_{l=0}^{N-1} \eta^{-l-1} \left(\partial_s^l K_M(t, s) \Big|_{s=0} - e^{-\eta t} \partial_s^l K_M(t, s) \Big|_{s=t} \right). \end{aligned}$$

This leads to (3.10) and completes the proof of Lemma 3.4. \square

The following Lemma 3.5 will be used in the proof of (ii) in Theorem 3.1.

Lemma 3.5. *Let $\{h_l\}_{l \in \mathbb{N}}$ and $\{p_l\}_{l \in \mathbb{N}}$ be given by (1.8). Then for each $l \in \mathbb{N}$,*

$$\partial_s^l K_M(t, s) \Big|_{s=0} = h_l(t) \quad \text{and} \quad \partial_s^l K_M(t, s) \Big|_{s=t} = -p_l(t), \quad t \geq 0. \quad (3.13)$$

Proof. Recall the conventional notation: $0^0 = 1$. First of all, we recall that \mathcal{M}_j is given in (2.4). Given $t \geq 0$, by (2.6), where $(\alpha, \beta) = (0, l)$, it follows that

$$\begin{aligned} \partial_s^l \mathcal{M}_j(t, s) \Big|_{s=0} &= \frac{(-1)^l}{j!} \sum_{m=0}^{\min\{l, j\}} C_l^m C_j^m m! \left((-s)^{j-m} \Big|_{s=0} \right) \left(\frac{d^{l-m}}{d\tau^{l-m}} \underbrace{M * \cdots * M}_j \right)(t) \\ &= \frac{(-1)^l}{j!} \left(\chi_{[0, l]}(j) C_l^j j! \right) \frac{d^{l-j}}{dt^{l-j}} \underbrace{M * \cdots * M}_j(t), \end{aligned}$$

which, along with (2.3), yields

$$\partial_s^l K_M(t, s) \Big|_{s=0} = \sum_{j=1}^{+\infty} \partial_s^l \mathcal{M}_j(t, s) \Big|_{s=0} = (-1)^l \sum_{j=0}^l C_l^j \frac{d^{l-j}}{dt^{l-j}} \underbrace{M * \cdots * M}_j(t).$$

This, along with (1.8), leads to the first equality in (3.13).

Next, from (2.6) with $(\alpha, \beta) = (0, l)$, it follows that

$$\partial_s^l \mathcal{M}_j(t, s) \Big|_{s=t} = \frac{(-1)^l}{j!} \sum_{m=0}^{\min\{l, j\}} C_l^m C_j^m m! (-t)^{j-m} \left(\frac{d^{l-m}}{d\tau^{l-m}} \underbrace{M * \cdots * M}_j \right)(0). \quad (3.14)$$

Meanwhile, by Lemma 2.1, we find

$$\frac{d^{l-m}}{d\tau^{l-m}} \underbrace{M * \cdots * M}_j(0) = 0, \quad \text{when } l - m < j - 1.$$

From the above and (3.14), we see

$$\partial_s^l \mathcal{M}_j(t, s) \Big|_{s=t} = (-1)^l \sum_{m=0}^{\min\{l, j\}} C_l^m \frac{(-t)^{j-m}}{(j-m)!} \chi_{[0, l-m]}(j-1) \frac{d^{l-m}}{d\tau^{l-m}} \underbrace{M * \cdots * M}_j(0).$$

This, along with (2.3), yields

$$\begin{aligned}\partial_s^l K_M(t, s)|_{s=t} &= \sum_{j=1}^{+\infty} \partial_s^l \mathcal{M}_j(t, s)|_{s=t} \\ &= (-1)^l \sum_{j=1}^{l+1} \sum_{m=0}^{\min\{l, j, l-j+1\}} \frac{(-t)^{j-m}}{(j-m)!} \left(C_l^{l-m} \frac{d^{l-m}}{d\tau^{l-m}} \underbrace{M * \cdots * M}_j(0) \right).\end{aligned}$$

Replacing $j - m$ by a new variable q in the above, using (1.8), we obtain the second equality in (3.13).

Hence, we finish the proof of Lemma 3.5. \square

We now are on the position to prove Theorem 3.1.

Proof of Theorem 3.1. Fix $N \geq 2$ and $\eta > 0$ arbitrarily. The proof is structured in three steps.

Step 1. Proof of (3.2).

By Lemmas 3.3, 3.4 and 3.5, and by (1.7), we find

$$\begin{aligned}w_\eta(t) &= e^{-\eta t} + e^{-\eta t} \sum_{l=0}^{N-1} \left(-\partial_s^l K_M(t, s)|_{s=t} \right) \eta^{-l-1} + \sum_{l=0}^{N-1} \left(\partial_s^l K_M(t, s)|_{s=0} \right) \eta^{-l-1} \\ &\quad + \eta^{-N} \int_0^t e^{-\eta s} \partial_s^N K_M(t, s) ds \\ &= e^{-\eta t} \sum_{l=0}^{N-1} \left(1 + p_l(t) \eta^{-l-1} \right) + \sum_{l=0}^{N-1} h_l(t) \eta^{-l-1} + \mathcal{R}_N(t, \eta) \eta^{-N-1}, \quad t \geq 0,\end{aligned}$$

which leads to (3.2).

Step 2. Proof of conclusion (i).

First, we have

$$\mathcal{R}_N \in C(\overline{\mathbb{R}^+} \times \overline{\mathbb{R}^+}). \quad (3.15)$$

Indeed, we apply Proposition 2.2 to see

$$K_M \in C^\infty(S_+). \quad (3.16)$$

From (1.7) and (3.16), (3.15) follows at once.

Second, we have

$$\mathcal{R}_N \in C(\mathbb{R}^+; C(\mathbb{R}^+)). \quad (3.17)$$

Indeed, it follows from (1.7) that when $t_2 \geq t_1 > 0$,

$$\begin{aligned}& \| \mathcal{R}_N(t_1, \cdot) - \mathcal{R}_N(t_2, \cdot) \|_{C(\mathbb{R}^+)} \\ & \leq \sup_{\tau > 0} \left[\int_0^{t_1} \tau e^{-\tau t} |\partial_s^N K_M(t_1, s) - \partial_s^N K_M(t_2, s)| ds + \int_{t_1}^{t_2} \tau e^{-\tau t} |\partial_s^N K_M(t_2, s)| ds \right] \\ & \leq \| \partial_s^N K_M(t_1, \cdot) - \partial_s^N K_M(t_2, \cdot) \|_{L^\infty(0, t_1)} + \frac{1}{t_1} \int_{t_1}^{t_2} |\partial_s^N K_M(t_2, s)| ds.\end{aligned}$$

This, along with (3.16), yields (3.17).

Third, from (1.7) and Proposition 2.2 (with $(\alpha, \beta) = (0, N)$), we see

$$\begin{aligned} \|\mathcal{R}_N(t, \cdot)\|_{C(\mathbb{R}^+)} &\leq \left(\sup_{\tau > 0} \int_0^\tau \tau e^{-\tau s} ds \right) \|\partial_s^N K_M(t, \cdot)\|_{L^\infty(0, t)} \\ &\leq e^t \left\{ \exp \left[N(1+t) \left(\sum_{j=0}^N \max_{0 \leq s \leq t} \left| \frac{d^j}{ds^j} M(s) \right| \right) \right] - 1 \right\}, \quad t \geq 0, \end{aligned}$$

which leads to (3.3).

Step 3. Proof of conclusion (ii).

Fix any $t \geq 0$. We apply Proposition 2.4 to $(t_0, s_0) = (t, 0)$ and $\vec{v} = (0, 1)$; $(t_0, s_0) = (t, t)$ and $\vec{v} = (0, 1)$, respectively, to find $j_1, j_2 \in \mathbb{N}$ so that

$$\partial_s^{j_1} K_M(t, 0) \neq 0 \quad \text{and} \quad \partial_s^{j_2} K_M(t, t) \neq 0. \quad (3.18)$$

Meanwhile, by (1.8) and (3.13), we have

$$h_0 \equiv 0, \quad \partial_s^{j_1} K_M(t, 0) = h_{j_1}(t) \quad \text{and} \quad \partial_s^{j_2} K_M(t, t) = -p_{j_2}(t). \quad (3.19)$$

Now, from (3.18) and (3.19), we see that neither $\{h_l(t)\}_{l \geq 1}$ nor $\{p_l(t)\}_{l \in \mathbb{N}}$ is the zero sequence.

This concludes the proof of Theorem 3.1. \square

4 Analysis of the memory-flow

In this section we present several technical propositions explaining the meaning of each component of the decomposition (1.9). We then prove Theorems 1.1-1.4 one by one. At last we prove some complementary properties of the flow and present an extension of Theorems 1.1-1.2.

4.1 On the components in the decomposition

The aim of this subsection is to explain the meaning of each component in the decomposition (1.9), and discuss their regularity properties. This will help us understanding the hybrid parabolic-hyperbolic behavior of the flow more deeply.

The next Proposition 4.1 (which is the conclusion (i) in Theorem 1.2) assures the smoothing effect of the first component in (1.9).

Proposition 4.1. *Let $N \geq 2$ be an integer. Then $\mathcal{P}_N(t)\mathcal{H}^{-\infty} \subset \mathcal{H}^{+\infty}$ for all $t > 0$.*

Proof. This directly follows from the definition of $\mathcal{P}_N(\cdot)$ (in (1.10)) and the smoothing effect of the heat semigroup $\{e^{tA}\}_{t \geq 0}$. \square

The next Proposition 4.2 (which contains the conclusion (ii) in Theorem 1.2) shows the propagation of singularities along the time direction for the second component $\mathcal{W}_N(\cdot)$. This shows the hyperbolic nature of $\mathcal{W}_N(\cdot)$, with null velocity of propagation.

Proposition 4.2. *Let $N \geq 2$ be an integer and let $y_0 \in \mathcal{H}^{-\infty}$ and $x_0 \in \Omega$. Then the following statements are equivalent:*

- (i) *For some $t_0 > 0$, $\mathcal{W}_N(\cdot)y_0 \notin L_{loc}^2(t_0, x_0)$;*
- (ii) *For each $t > 0$, $\mathcal{W}_N(\cdot)y_0 \notin L_{loc}^2(t, x_0)$;*
- (iii) *It holds that $A^{-2}y_0 \notin L_{loc}^2(x_0)$.*

Proof. We first claim that if $z \in \mathcal{H}^{-\infty}$ satisfies $z \in L_{loc}^2(x_0)$, then

$$A^{-1}z \in L_{loc}^2(x_0). \quad (4.1)$$

The proof of (4.1) is classical and, for the sake of completeness, we present it as follows: Write $g := A^{-1}z$. It is clear that $Ag \in L_{loc}^2(x_0)$. Then by (1.3) and some computations, we can see that $\Delta g \in L_{loc}^2(x_0)$. Since Δ is elliptic at x_0 , one has that $g|_{B(x_0, r)} \in H^2(B(x_0, r))$ for some $r > 0$ (see, for instance, [12, Theorem 18.1.29]). Thus, $g \in L_{loc}^2(x_0)$, which gives that $A^{-1}z \in L_{loc}^2(x_0)$.

We next prove that (i) \Rightarrow (iii). By contradiction, we suppose that (i) is true, but

$$A^{-2}y_0 \in L_{loc}^2(x_0). \quad (4.2)$$

Since M is analytic, it follows from (1.8) that $h_0 \equiv 0$ and each h_l is smooth. Then by (4.2) and (4.1), one has

$$\sum_{l=0}^{N-1} h_l(\cdot)(-A)^{-l-1}y_0 = \sum_{l=1}^{N-1} h_l(\cdot)(-A)^{-l+1}(A^{-2}y_0) \in L_{loc}^2(t, x_0) \quad \text{for each } t > 0.$$

This, along with the definition of $\mathcal{W}_N(\cdot)$ (in (1.10)), implies that

$$\mathcal{W}_N(\cdot)y_0 \in L_{loc}^2(t, x_0) \quad \text{for each } t > 0,$$

which contradicts (i). Therefore, (iii) is true.

We now show that (iii) \Rightarrow (ii). By contradiction, we suppose that (iii) holds, but $\mathcal{W}_N(\cdot)y_0 \in L_{loc}^2(\hat{t}_0, x_0)$ for some $\hat{t}_0 > 0$. Then from (1.10), we find

$$\sum_{l=0}^{N-1} h_l(\cdot)(-A)^{-l-1}y_0 \in L_{loc}^2(\hat{t}_0, x_0). \quad (4.3)$$

We will use (4.3) to prove that

$$A^{-2}y_0 \in L_{loc}^2(x_0). \quad (4.4)$$

When this is done, we are led to a contradiction with (iii), and then the statement (ii) is true.

To prove (4.4), we observe that there are only two possibilities: either $N = 2$ or $N \geq 3$. When that $N = 2$, we see from (4.3) and (1.8) that

$$-M(\cdot)A^{-2}y_0 = \sum_{l=0}^{N-1} h_l(\cdot)(-A)^{-l-1}y_0 \in L_{loc}^2(\hat{t}_0, x_0).$$

Since M is nonzero, (4.4) (for this case) follows by integrating the above in the time variable.

When $N \geq 3$, it follows from (1.8) that $h_0 = 0$ and $h_1 = -M$. Thus, the leading term of the sum in (4.3) is $-M(\cdot)A^{-2}$. From this and the fact that M is nonzero, by integrating (4.3) in the time variable, we see

$$A^{-2}y_0 + \sum_{l=2}^{N-1} c_l A^{-l-1}y_0 \in L_{loc}^2(x_0) \quad \text{for some } \{c_l\}_{l=2}^{N-1} \subset \mathbb{R}.$$

Write $\mathcal{A} := -\sum_{l=2}^{N-1} c_l A^{-l+1}$. Then the above leads to

$$f_0 := (1 - \mathcal{A})(A^{-2}y_0) \in L_{loc}^2(x_0). \quad (4.5)$$

Since $y_0 \in \mathcal{H}^{-\infty}$, there is $m \in \mathbb{N}^+$ so that $y_0 \in \mathcal{H}^{-m}$. This, along with (4.5), yields

$$A^{-2}y_0 - (1 + \mathcal{A} + \cdots + \mathcal{A}^{m-1})f_0 = \mathcal{A}^m(A^{-2}y_0) \in L^2(\Omega). \quad (4.6)$$

At the same time, since $f_0 \in L_{loc}^2(x_0)$ in (4.5), it follows from (4.1) that

$$\mathcal{A}^j f_0 \in L_{loc}^2(x_0) \quad \text{for each } j \in \mathbb{N}.$$

Thus, (4.4) (for $N \geq 3$) follows from (4.6). Hence, (4.4) is true.

Finally, it is clear that (ii) \Rightarrow (i). Hence, we finish the proof of Proposition 4.2. \square

The following Proposition 4.3 (which is partially the conclusion (iii) in Theorem 1.2) gives an important time-uniform smoothing effect of the last component $\mathfrak{R}_N(\cdot)$.

Proposition 4.3. *Let $N \geq 2$ be an integer. Then the last component \mathfrak{R}_N in (1.9) exhibits a time-uniform smoothing effect with a gain of $2N + 2$ space derivatives: for each $y_0 \in \mathcal{H}^s$ with $s \in \mathbb{R}$,*

$$\mathfrak{R}_N(\cdot)y_0 \in C(\overline{\mathbb{R}^+}; \mathcal{H}^{s+2N+2}), \quad \text{while } A^{-j}e^{\cdot A}y_0, A^{-j}y_0 \in C(\overline{\mathbb{R}^+}; \mathcal{H}^{s+2j}) \quad (0 \leq j \leq N). \quad (4.7)$$

Remark 4.4. There is a delicate point worth to clarifying. Although the map $t \mapsto A^{-j}e^{tA}$, $t \geq 0$ (for a fixed j) has infinite order smoothing effect at positive time, it has the finite time-uniform smoothing effect (given in (4.7)) with only $2j$ space-derivatives gained. This index $2j$ is optimal due to the following fact: for given $y_0 \in \mathcal{H}^{-\infty}$ and $s \in \mathbb{R}$, $A^{-j}e^{\cdot A}y_0 \in C(\overline{\mathbb{R}^+}; \mathcal{H}^{s+2j})$ if and only if $y_0 \in \mathcal{H}^s$, which can be directly checked.

Proof of Proposition 4.3. The second statement in (4.7) is clearly true. We now show the first statement in (4.7). To this end, we arbitrarily fix $s \in \mathbb{R}$, $y_0 \in \mathcal{H}^s$ and $t_0 \geq 0$. We aim to show the continuity of $\mathfrak{R}_N(\cdot)y_0$ at time t_0 . Fix any $\varepsilon > 0$. Write $y_0 = \sum_{j=1}^{\infty} y_{0,j}e_j$. Since $y_0 \in \mathcal{H}^s$, we can choose $j_\varepsilon \in \mathbb{N}^+$ large enough so that $\sum_{j \geq j_\varepsilon} \eta_j^s y_{0,j}^2 < \varepsilon^2$. This, along with the definitions of $\mathfrak{R}_N(\cdot)$ (see (1.10)) and \mathcal{R}_N (see (1.7)), yields that for each $t \in [0, t_0 + 1]$,

$$\begin{aligned} & \|\mathfrak{R}_N(t)y_0 - \mathfrak{R}_N(t_0)y_0\|_{\mathcal{H}^{s+2N+2}}^2 \\ &= \|\mathcal{R}_N(t, -A)y_0 - \mathcal{R}_N(t_0, -A)y_0\|_{\mathcal{H}^s}^2 \\ &= \sum_{j \geq 1} |\mathcal{R}_N(t, \eta_j) - \mathcal{R}_N(t_0, \eta_j)|^2 y_{0,j}^2 \eta_j^s \\ &\leq \max_{1 \leq j \leq j_\varepsilon} |\mathcal{R}_N(t, \eta_j) - \mathcal{R}_N(t_0, \eta_j)|^2 \left(\sum_{j \geq 1} y_{0,j}^2 \eta_j^s \right) + 4\|\mathcal{R}_N\|_{C([0, t_0+1] \times \mathbb{R}^+)}^2 \varepsilon^2. \end{aligned}$$

From this and (3.3), one can find some $C > 0$ (independent of ε) and $\delta \in (0, 1)$ so that

$$\|\mathfrak{R}_N(t)y_0 - \mathfrak{R}_N(t_0)y_0\|_{\mathcal{H}^{s+2N+2}}^2 \leq C\varepsilon^2, \quad \text{when } t \in (t_0 - \delta, t_0 + \delta) \cap \mathbb{R}^+.$$

This leads to the continuity of $\mathfrak{R}_N(\cdot)y_0$ at time t_0 and ends the proof of Proposition 4.3. \square

4.2 Proof of main theorems

We now give the proofs of Theorems 1.1-1.4 one by one.

Proof of Theorem 1.1. First of all, (1.11) (with $t > 0$) follows from the conclusion (ii) of Theorem 3.1 at once.

The remainder is to show (1.9). Fix $s \in \mathbb{R}$ and $y_0 \in \mathcal{H}^s$. Recall that η_j is the j^{th} eigenvalue of $-A$ and e_j is the corresponding normalized eigenfunction in $L^2(\Omega)$. Then we can write

$$y_0 = \sum_{j \geq 1} y_{0,j}e_j \quad \text{and} \quad y(t; y_0) = \sum_{j \geq 1} y_j(t)e_j, \quad t \geq 0, \quad (4.8)$$

where $y_{0,j} := \langle y_0, e_j \rangle_{\mathcal{H}^s, \mathcal{H}^{-s}}$ and $y_j(\cdot)$ satisfies

$$y_j'(t) + \eta_j y_j(t) + \int_0^t M(t-\tau) y_j(\tau) d\tau = 0, \quad t > 0; \quad y_j(0) = y_{0,j}. \quad (4.9)$$

By (4.9) and (3.1), we see that for each $j \in \mathbb{N}^+$,

$$y_j(t) = y_{0,j} w_{\eta_j}(t), \quad t \geq 0, \quad (4.10)$$

where $w_{\eta_j}(\cdot)$ is the solution to (3.1) with $\eta = \eta_j$. Then by (4.8) and (4.10), we have

$$y(t; y_0) = \sum_{j \geq 1} w_{\eta_j}(t) y_{0,j} e_j, \quad t \geq 0, \quad (4.11)$$

From (4.11) and (3.2) (in Theorem 3.1), we see that when $t \geq 0$,

$$y(t; y_0) = \sum_{j \geq 1} \left[e^{-\eta_j t} \left(1 + \sum_{l=0}^{N-1} p_l(t) \eta_j^{-l-1} \right) + \sum_{l=0}^{N-1} h_l(t) \eta_j^{-l-1} + \mathcal{R}_N(t, \eta_j) \eta_j^{-N-1} \right] y_{0,j} e_j. \quad (4.12)$$

Meanwhile, by functional calculus, we have

$$\text{RHS of (4.12)} = \left[\left(e^{tA} + e^{tA} \sum_{l=0}^{N-1} p_l(t) (-A)^{-l-1} \right) + \sum_{l=0}^{N-1} h_l(t) (-A)^{-l-1} + \mathcal{R}_N(t, -A) (-A)^{-N-1} \right] y_0. \quad (4.13)$$

Here, RHS of (4.12) denotes the expression on the right-hand side of (4.12).

Since y_0 was arbitrarily taken from \mathcal{H}^s , we can use (4.12), (4.13), (1.10) and Proposition 7.1 (in Appendix) to get (1.9). This concludes the proof of Theorem 1.1. \square

Proof of Theorem 1.2. First of all, the conclusions (i)-(ii) in Theorem 1.2 follow from Proposition 4.1 and Proposition 4.2, respectively. In what follows, the conclusions (iii)-(iv) in Theorem 1.2 will be proved one by one.

Step 1. The proof of the conclusion (iii) in Theorem 1.2

By Proposition 4.3, we only need to show (1.13). For this purpose, we first claim

$$\mathcal{R}_N(\cdot, -A)|_{\mathbb{R}^+} \in C(\mathbb{R}^+; \mathcal{L}(\mathcal{H}^s)) \text{ for each } s \in \mathbb{R}. \quad (4.14)$$

To prove (4.14), we arbitrarily fix $s \in \mathbb{R}$ and $t_2 \geq t_1 > 0$. Set

$$\tilde{\mathcal{R}}(-A) := \mathcal{R}_N(t_2, -A) - \mathcal{R}_N(t_1, -A). \quad (4.15)$$

Then $\tilde{\mathcal{R}}(-A)$ is the operator obtained by the functional calculus of the function: $\mathcal{R}_N(t_2, \cdot) - \mathcal{R}_N(t_1, \cdot)$. By (1.4) and by the spectral representation of $\tilde{\mathcal{R}}(-A)$, we see

$$\|\tilde{\mathcal{R}}(-A)\|_{\mathcal{L}(\mathcal{H}^s)}^2 = \sup_{\sum \eta_j^s z_j^2 \leq 1} \left\| \sum_{j \geq 1} \tilde{\mathcal{R}}(\eta_j) z_j e_j \right\|_{\mathcal{H}^s}^2 = \sup_{\sum \eta_j^s z_j^2 \leq 1} \sum_{j=1}^{+\infty} \eta_j^s |\tilde{\mathcal{R}}(\eta_j)|^2 z_j^2 \leq \sup_{\eta > 0} |\tilde{\mathcal{R}}(\eta)|^2.$$

This, along with (4.15), yields

$$\|\mathcal{R}_N(t_2, -A) - \mathcal{R}_N(t_1, -A)\|_{\mathcal{L}(\mathcal{H}^s)} = \|\tilde{\mathcal{R}}(-A)\|_{\mathcal{L}(\mathcal{H}^s)} \leq \|\mathcal{R}_N(t_2, \cdot) - \mathcal{R}_N(t_1, \cdot)\|_{C(\mathbb{R}^+)}. \quad (4.16)$$

Since $\mathcal{R}_N \in C(\mathbb{R}^+; C(\mathbb{R}^+))$ (see Theorem 3.1), (4.14) follows from (4.16) at once.

We now show (1.13). Indeed, arguing as in the proof of (4.16), we can obtain

$$\|\mathcal{R}_N(t, -A)\|_{\mathcal{L}(\mathcal{H}^s)} \leq \|\mathcal{R}_N(t, \cdot)\|_{C(\mathbb{R}^+)}, \quad t \geq 0.$$

This, along with (3.3) (in Theorem 3.1), leads to (1.13).

Hence, the conclusion (iii) in Theorem 1.2 is true.

Step 2. The proof of the conclusion (iv) in Theorem 1.2

Arbitrarily fix $y_0 \in \mathcal{H}^{-\infty}$, $x_0 \in \Omega$ and $t > 0$. There is an integer $m \geq 2$ so that

$$y_0 \in \mathcal{H}^{-2m}.$$

Then by (1.9) and (1.13) (where N is replaced by m), one can easily check

$$\Phi(\cdot)y_0 - \mathcal{P}_m(\cdot)y_0 - \mathcal{W}_m(\cdot)y_0 \in L_{loc}^\infty(\overline{\mathbb{R}^+}; L^2(\Omega)). \quad (4.17)$$

Combine (4.17) and Proposition 4.1 (where N is replaced by m) to get

$$\Phi(\cdot)y_0 \notin L_{loc}^2(t, x_0) \Leftrightarrow \mathcal{W}_m(\cdot)y_0 \notin L_{loc}^2(t, x_0). \quad (4.18)$$

Meanwhile, for each integer $k \geq 2$, it follows by Proposition 4.2 (where N is replaced by k) that

$$\mathcal{W}_k(\cdot)y_0 \notin L_{loc}^2(t, x_0) \Leftrightarrow A^{-2}y_0 \notin L_{loc}^2(x_0),$$

in particular,

$$\mathcal{W}_m(\cdot)y_0 \notin L_{loc}^2(t, x_0) \Leftrightarrow \mathcal{W}_N(\cdot)y_0 \notin L_{loc}^2(t, x_0).$$

This, along with (4.18), yields

$$\Phi(\cdot)y_0 \notin L_{loc}^2(t, x_0) \Leftrightarrow \mathcal{W}_N(\cdot)y_0 \notin L_{loc}^2(t, x_0). \quad (4.19)$$

Thus, the conclusion (iv) follows from (4.19) and Proposition 4.2 at once.

Hence, we complete the proof of Theorem 1.2. \square

Proof of Theorem 1.3. Arbitrarily fix an integer $N \geq 2$. We first show (1.15). Indeed, one can see from (1.2), (1.10) and (1.7) that for each $j \in \mathbb{N}^+$,

$$\begin{aligned} \Phi(0)e_j &= e_j, \quad \mathcal{P}_N(0)e_j = \left(1 + \sum_{l=0}^{N-1} p_l(0)\eta_j^{-l-1}\right)e_j, \\ \mathfrak{R}_N(0)e_j &= 0, \quad \mathcal{W}_N(0)e_j = \left(\sum_{l=0}^{N-1} h_l(0)\eta_j^{-l-1}\right)e_j. \end{aligned}$$

Since $\lim_{j \rightarrow +\infty} \eta_j = +\infty$, the above leads to (1.15) at once.

We next prove (1.16). Arbitrarily fix $t > 0$. On one hand, it follows from (1.8) that $h_0 \equiv 0$ and $h_1(t) = -M(t)$, which, along with (1.10), yield

$$\mathcal{W}_N(t)e_j = -M(t)\eta_j^{-2}e_j + \sum_{1 \leq l \leq N-1} h_l(t)\eta_j^{-l-1}e_j \quad \text{for all } j \geq 1.$$

This implies

$$\lim_{j \rightarrow +\infty} \|\mathcal{W}_N(t)e_j\|_{\mathcal{H}^4} = |M(t)| \quad \text{and} \quad \lim_{j \rightarrow +\infty} \|\mathcal{W}_N(t)e_j\|_{\mathcal{H}^s} = 0, \quad \forall s < 4. \quad (4.20)$$

On the other hand, from (1.10), it follows that for each $j \geq 1$,

$$\mathcal{P}_N(t)e_j = e^{-t\eta_j} \left(1 + \sum_{l=0}^{N-1} p_l(t)\eta_j^{-l-1}\right)e_j \quad \text{and} \quad \mathfrak{R}_N(t)e_j = \eta_j^{-N-1}\mathcal{R}_N(t, -A)e_j.$$

Since $\lim_{j \rightarrow +\infty} \eta_j^{-1} = \lim_{j \rightarrow +\infty} \eta_j^{\frac{s}{2}} e^{-t\eta_j} = 0$ ($s \in \mathbb{R}$), the above, together with (1.13), gives

$$\lim_{j \rightarrow +\infty} \|\mathcal{P}_N(t)e_j\|_{\mathcal{H}^{s_1}} = \lim_{j \rightarrow +\infty} \|\mathfrak{R}_N(t)e_j\|_{\mathcal{H}^{s_2}} = 0, \quad s_1 \in \mathbb{R}, \quad s_2 < 2N + 2. \quad (4.21)$$

Now, by (4.20) and (4.21), we can use the decomposition (1.9) (with the above $N \geq 2$) to get

$$\lim_{j \rightarrow +\infty} \|\Phi(t)e_j\|_{\mathcal{H}^4} = |M(t)| \quad \text{and} \quad \lim_{j \rightarrow +\infty} \|\Phi(t)e_j\|_{\mathcal{H}^s} = 0, \quad \forall s < 4.$$

These, along with (4.20)-(4.21), lead to (1.16). This completes the proof of Theorem 1.3. \square

Proof of Theorem 1.4. Arbitrarily fix $s \in \mathbb{R}$. We will prove the conclusions (i)-(iii) one by one.

(i) Arbitrarily fix $\alpha \in [0, 4]$ and $t > 0$. We apply (1.9) (with $N = 2$), as well as (1.10) and (1.8), to obtain

$$\Phi(t) = \left(e^{tA} - p_0(t)A^{-1}e^{tA} + p_1(t)A^{-2}e^{tA} \right) - M(t)A^{-2} + \mathcal{R}_2(t, -A)(-A)^{-3}. \quad (4.22)$$

Meanwhile, notice that

$$\|A^{\frac{\alpha}{2}-j}e^{tA}\|_{\mathcal{L}(\mathcal{H}^s)} \left(= \sup_{j \in \mathbb{N}^+} \eta_j^{\frac{\alpha}{2}-j} e^{-t\eta_j} \right) \leq t^{j-\frac{\alpha}{2}} \quad \text{and} \quad \|A^{\frac{\alpha}{2}-j-2}\|_{\mathcal{L}(\mathcal{H}^s)} \leq \eta_1^{\frac{\alpha}{2}-j-2}, \quad j = 0, 1, 2.$$

These, along with (4.22), yield

$$\|\Phi(t)\|_{\mathcal{L}(\mathcal{H}^s, \mathcal{H}^{s+\alpha})} \leq t^{-\frac{\alpha}{2}} \left(1 + t|p_0(t)| + t^2|p_1(t)| \right) + \eta_1^{\frac{\alpha}{2}-2}|M(t)| + \eta_1^{\frac{\alpha}{2}-3}\|\mathcal{R}_2(t, \cdot)\|_{\mathcal{L}(\mathcal{H}^{s+\alpha})}.$$

After direct computations, we obtain (1.17) from the above, (1.8) and (1.13) (where $N = 2$).

(ii) Assume that (1.18) holds for $t \in (0, \delta_0)$. We first claim

$$+\infty > \limsup_{j \rightarrow +\infty} \eta_j^{\frac{\alpha_0}{2}-2}, \quad \text{i.e.,} \quad \alpha_0 \leq 4. \quad (4.23)$$

Indeed, by the assumption (C), we can choose small $t_0 \in (0, \delta_0)$ so that

$$M(t_0) \neq 0. \quad (4.24)$$

Then from (4.22) and (1.13) (where $N = 2$), it follows that for each $j \geq 1$,

$$\|\Phi(t_0)\|_{\mathcal{L}(\mathcal{H}^s, \mathcal{H}^{s+\alpha_0})} \geq \|\Phi(t_0)e_j\|_{\mathcal{H}^{s+\alpha_0}} / \|e_j\|_{\mathcal{H}^s} = \eta_j^{\frac{\alpha_0}{2}-2} \left(|M(t_0)| + O(\eta_j^{-1}) \right),$$

(Here and what follows, by $O(\eta_j^{-1})$ with $j \geq 1$, we mean that there is $C_1 > 0$, independent of j , so that $|O(\eta_j^{-1})| \leq C_1 \eta_j^{-1}$.) This, along with (1.18) (where $t = t_0$), (4.24) and the fact that $\lim_{j \rightarrow +\infty} \eta_j = +\infty$, leads to (4.23).

We next claim

$$\liminf_{t \rightarrow 0^+} t^{\frac{\alpha_0}{2}} \|\Phi(t)\|_{\mathcal{L}(\mathcal{H}^s, \mathcal{H}^{s+\alpha_0})} > 0. \quad (4.25)$$

In fact, from (1.18) (where $z = e_j$), (4.22) and (1.13), we see that when $t \in (0, \delta_0)$ and $j \in \mathbb{N}^+$,

$$\begin{aligned} t^{\frac{\alpha_0}{2}} \|\Phi(t)\|_{\mathcal{L}(\mathcal{H}^s, \mathcal{H}^{s+\alpha_0})} &\geq t^{\frac{\alpha_0}{2}} \|\Phi(t)e_j\|_{\mathcal{H}^{s+\alpha_0}} / \|e_j\|_{\mathcal{H}^s} \\ &\geq (t\eta_j)^{\frac{\alpha_0}{2}} \left(|1 + O(\eta_j^{-1})| e^{-t\eta_j} - O(\eta_j^{-1}) \right). \end{aligned} \quad (4.26)$$

Meanwhile, it follows by Weyl's asymptotic formula for the eigenvalues (see for instance [16, XIII.15]) that $\lim_{j \rightarrow +\infty} \eta_{j+1}/\eta_j = 1$. Thus, there is $j_0 \in \mathbb{N}^+$ so that

$$[\eta_{j_0}, +\infty) = \cup_{j \geq j_0} [\eta_j, 2\eta_j).$$

Therefore, for each $t \in (0, \eta_{j_0}^{-1}]$, there is an integer $j_t \geq j_0$ so that

$$t^{-1} \in [\eta_{j_t}, 2\eta_{j_t}), \text{ i.e., } \frac{1}{2} < t\eta_{j_t} \leq 1.$$

This, along with (4.26), leads to (4.25).

Finally, the conclusion (ii) follows from (4.23) and (4.25) at once.

(iii) We first claim

$$\Phi(\cdot)y_0 \in C(\mathbb{R}^+; \mathcal{H}^{s+4}) \text{ for each } y_0 \in \mathcal{H}^s. \quad (4.27)$$

To this end, we arbitrarily fix $y_0 \in \mathcal{H}^s$, $t_0 > 0$ and $\varepsilon > 0$. Since $y_0 \in \mathcal{H}^s$, there is $j_\varepsilon \in \mathbb{N}^+$ so that

$$\|y_0 - y_{0,j_\varepsilon}\|_{\mathcal{H}^s} < \varepsilon \text{ where } y_{0,j_\varepsilon} := \sum_{j=1}^{j_\varepsilon} \langle y_0, e_j \rangle_{\mathcal{H}^s, \mathcal{H}^{-s}} e_j. \quad (4.28)$$

At the same time, one can easily check $\Phi(\cdot)y_{0,j_\varepsilon} \in C(\mathbb{R}^+; \mathcal{H}^{s+4})$, which implies there is $\delta_\varepsilon \in (0, t_0/2)$ so that

$$\sup_{t_0 - \delta_\varepsilon < t < t_0 + \delta_\varepsilon} \|\Phi(t)y_{0,j_\varepsilon} - \Phi(t_0)y_{0,j_\varepsilon}\|_{\mathcal{H}^{s+4}} < \varepsilon.$$

This, together with (4.28), gives

$$\sup_{t_0 - \delta_\varepsilon < t < t_0 + \delta_\varepsilon} \|\Phi(t)y_0 - \Phi(t_0)y_0\|_{\mathcal{H}^{s+4}} < \varepsilon + 2 \sup_{t_0/2 \leq \tau \leq 2t_0} \|\Phi(\tau)\|_{\mathcal{L}(\mathcal{H}^s, \mathcal{H}^{s+4})} \varepsilon,$$

which, along with (1.17), leads to (4.27).

We next claim that for each $\alpha > 4$,

$$\Phi(\cdot)\hat{y}_0 \notin C(\mathbb{R}^+; \mathcal{H}^{s+\alpha}) \text{ for some } \hat{y}_0 \in \mathcal{H}^s. \quad (4.29)$$

For this purpose, we arbitrarily fix $\alpha > 4$. Since $h_0(t) \equiv 0$ and $h_1(t) = -M(t)$ (see (1.8)), we apply (1.9) (with $N = 2$), as well as (1.10) and (1.13), to obtain

$$\lim_{j \rightarrow +\infty} \eta_j^2 \langle \Phi(t)e_j, e_j \rangle_{L^2(\Omega)} = -M(t) \text{ for each } t > 0. \quad (4.30)$$

Meanwhile, by the assumption (\mathfrak{C}) , we can choose $\hat{t}_0 > 0$ so that $M(\hat{t}_0) \neq 0$. This, along with (4.30) and the fact that $\alpha > 4$, yields

$$\eta_j^{\frac{\alpha}{2}} |\langle \Phi(\hat{t}_0) e_j, e_j \rangle_{L^2(\Omega)}| \rightarrow +\infty \text{ as } j \rightarrow +\infty.$$

Thus we can choose a subsequence $\{k_j\}_{j \geq 1}$ of \mathbb{N}^+ so that

$$\eta_{k_j}^{\frac{\alpha}{2}} |\langle \Phi(\hat{t}_0) e_{k_j}, e_{k_j} \rangle_{L^2(\Omega)}| \geq j 2^j \text{ for each } j \geq 1. \quad (4.31)$$

We now define $\hat{y}_0 \in \mathcal{H}^s$ as follows:

$$\hat{y}_0 := \sum_{j \geq 1} 2^{-j} \eta_{k_j}^{-\frac{s}{2}} e_{k_j}. \quad (4.32)$$

Then, by (4.31) and (4.32), we find

$$\begin{aligned} \|\Phi(\hat{t}_0) \hat{y}_0\|_{\mathcal{H}^{s+\alpha}}^2 &= \sum_{j \geq 1} \eta_{k_j}^{s+\alpha} \left(\langle \Phi(\hat{t}_0) e_{k_j}, e_{k_j} \rangle_{L^2(\Omega)} \langle \hat{y}_0, e_{k_j} \rangle_{\mathcal{H}^s, \mathcal{H}^{-s}} \right)^2 \\ &= \sum_{j \geq 1} \left(\eta_{k_j}^{\frac{\alpha}{2}} \langle \Phi(\hat{t}_0) e_{k_j}, e_{k_j} \rangle_{L^2(\Omega)} 2^{-j} \right)^2 \geq \sum_{j \geq 1} j^2 = +\infty, \end{aligned}$$

which implies $\Phi(\hat{t}_0) \hat{y}_0 \notin \mathcal{H}^{s+\alpha}$. This leads to (4.29).

In conclusion, we complete the proof of Theorem 1.4. \square

Remark 4.5. By a very similar way to that used in the proof of (1.17) (in Theorem 1.4), we can also show what follows: For each $t > 0$,

$$\Phi(t) \in \mathcal{L}(\mathcal{H}^s, \mathcal{H}^{s+2k(t)+2}) \text{ for each } s \in \mathbb{R}, \quad (4.33)$$

where $k(t) := \min\{l \geq 1 : h_l(t) \neq 0\}$.

From (4.33) and the fact that $h_1(t) = -M(t)$ (which follows from (1.8)), we conclude that *the smoothing effect of the flow $\Phi(t)$ at points in the set $\{t > 0 : M(t) = 0\}$ is better than that at points in the set $\{t > 0 : M(t) \neq 0\}$.*

4.3 Other properties of the flow and the components

This subsection presents more properties of the flow and the components. More precisely, first, we formulate Propositions 4.6 to illustrate how the component \mathcal{P}_N influences the flow, from the perspective of the singularities; (In plain language, it tells us that the singularity of $\mathcal{P}_N(0)$ determines the singularity of $\Phi(0)$.); second, we give Proposition 4.7 to show that both $\mathcal{P}_N(0)$ and $\mathcal{W}_N(0)$ are not projections; last, we present Propositions 4.8-4.9, which might have independent interest.

Proposition 4.6. *Let $N \geq 2$ be an integer and let $y_0 \in \mathcal{H}^{-\infty}$ and $x_0 \in \Omega$. Then $(y_0 =) \Phi(0)y_0 \notin L_{loc}^2(x_0)$ if and only if $\mathcal{P}_N(0)y_0 \notin L_{loc}^2(x_0)$.*

Proof. First of all, since $\Phi(0)y_0 = y_0$, we have

$$\Phi(0)y_0 \notin L_{loc}^2(x_0) \Leftrightarrow y_0 \notin L_{loc}^2(x_0). \quad (4.34)$$

We claim

$$\mathcal{P}_N(0)y_0 \notin L_{loc}^2(x_0) \Leftrightarrow y_0 \notin L_{loc}^2(x_0) \quad (4.35)$$

When this is proved, the conclusion in this proposition follows from (4.34) and (4.35) at once.

We now show (4.35). To prove the necessity, we suppose, by contradiction, that the statement on the left-hand side of (4.35) is true, but $y_0 \in L_{loc}^2(x_0)$. Then we apply (4.1) to obtain that

$$A^{-j}y_0 \in L_{loc}^2(x_0) \quad \text{for each } j \in \mathbb{N}.$$

At the same time, it follows from (1.10) that

$$\mathcal{P}_N(0)y_0 = y_0 + \sum_{l=0}^{N-1} p_l(0)(-A)^{-l-1}y_0.$$

These imply that $\mathcal{P}_N(0)y_0 \in L_{loc}^2(x_0)$, which contradicts the statement on the left-hand side of (4.35). Therefore, we have shown the necessity.

To show the sufficiency, we suppose, by contradiction, that the statement on the right-hand side of (4.35) holds, but $\mathcal{P}_N(0)y_0 \in L_{loc}^2(x_0)$. Then from (1.10), there is a sequence $(\hat{c}_l)_{l=0}^{N-1} \subset \mathbb{R}$ so that

$$y_0 + \sum_{l=0}^{N-1} \hat{c}_l A^{-l-1}y_0 \in L_{loc}^2(x_0).$$

Then by the similar way as that used in the proof of “(4.3) \Rightarrow (4.4)”, one can get that $y_0 \in L_{loc}^2(x_0)$. This contradicts the statement on the right-hand side of (4.35). Therefore, the sufficiency is proved. Hence, we finish the proof of Proposition 4.6. \square

Proposition 4.7. *The following conclusions are true:*

(i) *For each integer $N \geq 2$,*

$$\mathfrak{R}_N(0) = 0 \quad \text{and} \quad y_0 = \mathcal{P}_N(0)y_0 + \mathcal{W}_N(0)y_0 \quad \text{for each } y_0 \in L^2(\Omega). \quad (4.36)$$

(ii) *For each integer $N \geq 2$, $\mathcal{P}_N(0)$ and $\mathcal{W}_N(0)$ are projections over $L^2(\Omega)$ if and only if*

$$M(0) = \cdots = M^{(j)}(0) = \cdots = M^{(N-2)}(0) = 0. \quad (4.37)$$

Proof. First, the conclusion (i) follows from (1.10), (1.7) and (1.9) (with $t = 0$) at once.

We next prove the conclusion (ii). To show the sufficiency, we assume that (4.37) is true. Then by (4.37), we can apply (2.1) (with $t \rightarrow 0^+$) to get

$$\frac{d^l}{dt^l} \underbrace{M * \cdots * M(0)}_j = 0, \quad j \in \mathbb{N}^+, \quad l \in \{0, \dots, N-2\}.$$

This, along with (1.8), yields

$$h_l(0) = 0, \quad 0 \leq l \leq N-1,$$

from which and (1.10), it follows $\mathcal{W}_N(0) = 0$ and consequently, $\mathcal{W}_N(0)$ is a projection. Then by the second equality in (4.36), we see that $\mathcal{P}_N(0)$ is a projection.

To prove the necessity, we suppose that $\mathcal{P}_N(0)$ and $\mathcal{W}_N(0)$ are projections over $L^2(\Omega)$. Then we have

$$\mathcal{W}_N(0)^2 y_0 = \mathcal{W}_N(0) y_0 \quad \text{for each } y_0 \in L^2(\Omega).$$

Taking $y_0 = e_j$ with $j \in \mathbb{N}^+$ in the above and using (1.10), we find

$$\begin{aligned} \left(\sum_{l=0}^{N-1} h_l(0) \eta_j^{-l-1} \right)^2 e_j &= \left(\sum_{l=0}^{N-1} h_l(0) (-A)^{-l-1} \right)^2 e_j = \mathcal{W}_N(0)^2 y_0 \\ &= \mathcal{W}_N(0) y_0 = \left(\sum_{l=0}^{N-1} h_l(0) (-A)^{-l-1} \right) e_j = \left(\sum_{l=0}^{N-1} h_l(0) \eta_j^{-l-1} \right) e_j. \end{aligned}$$

This implies

$$\left(\sum_{l=0}^{N-1} h_l(0) \eta_j^{-l-1} \right)^2 = \sum_{l=0}^{N-1} h_l(0) \eta_j^{-l-1} \quad \text{for all } j \geq 1.$$

Since $\lim_{j \rightarrow +\infty} \eta_j = +\infty$, the above, divided by $\eta_j^{-1}, \dots, \eta_j^{-N}$ respectively, gives $h_l(0) = 0$, $0 \leq l \leq N-1$. Then by direct computations and by (1.8), we get (4.37). This ends the proof of Proposition 4.7. \square

Proposition 4.8. *Let K_M be given by (1.6). Then*

$$\Phi(t)^* = \Phi(t) = e^{tA} + \int_0^t K_M(t, \tau) e^{\tau A} d\tau, \quad t \geq 0. \quad (4.38)$$

Moreover, it holds that

$$\sup_{s \in \mathbb{R}} \|\Phi(t) - e^{tA}\|_{\mathcal{L}(\mathcal{H}^s)} \leq \inf_{\lambda \geq -\eta_1} e^{\lambda t} \left[\exp \left(t \int_0^t e^{-\lambda \tau} |M(\tau)| d\tau \right) - 1 \right], \quad t \geq 0. \quad (4.39)$$

(Here, $-\eta_1$ is the first eigenvalue of A .)

Proof. Arbitrarily fix $s \in \mathbb{R}$ and $y_0 \in \mathcal{H}^s$. By (4.8), (4.11) and Lemma 3.3, we find

$$y(t; y_0) = \sum_{j \geq 1} w_{\eta_j}(t) y_{0,j} e_j = \sum_{j \geq 1} \left(e^{-\eta_j t} + \int_0^t K_M(t, \tau) e^{-\eta_j \tau} d\tau \right) y_{0,j} e_j, \quad t \geq 0. \quad (4.40)$$

Now the second equality in (4.38) follows from (4.40) and (1.2), while the first equality in (4.38) follows by the second one in (4.38) and the fact that for each $t \geq 0$, $e^{tA} = e^{tA^*}$ in $\mathcal{L}(\mathcal{H}^s)$.

Finally, it follows from (4.38) that

$$\|\Phi(t) - e^{tA}\|_{\mathcal{L}(\mathcal{H}^s)} \leq \int_0^t \|e^{\tau A}\|_{\mathcal{L}(\mathcal{H}^s)} |K_M(t, \tau)| d\tau \leq \int_0^t e^{-\eta_1 \tau} |K_M(t, \tau)| d\tau, \quad t \geq 0.$$

This, along with (2.14) in Proposition 2.6, where $\lambda \geq -\eta_1$, yields

$$\|\Phi(t) - e^{tA}\|_{\mathcal{L}(\mathcal{H}^s)} \leq e^{\lambda t} \int_0^t e^{-\lambda(t-\tau)} |K_M(t, \tau)| d\tau \leq e^{\lambda t} \left[\exp \left(t \int_0^t e^{-\lambda \tau} |M(\tau)| d\tau \right) - 1 \right], \quad t \geq 0,$$

which leads to (4.39) and completes the proof of Proposition 4.8. \square

Proposition 4.9. *Let $s \in \mathbb{R}$. Then $\Phi(\cdot)$ is real analytic from \mathbb{R}^+ to $\mathcal{L}(\mathcal{H}^s)$.*

Proof. According to Proposition 2.3, K_M is real analytic over S_+ . Then by (4.38) and the analyticity of $\{e^{tA}\}_{t>0}$, we obtain that $\Phi(\cdot)$ is real analytic from \mathbb{R}^+ to $\mathcal{L}(\mathcal{H}^s)$. This ends the proof of Proposition 4.9. \square

4.4 Less regular memory kernels

The techniques of this paper can also be employed to handle less regular memory kernels. Assume that:

(\mathfrak{C}_1) The memory kernel M is in $C^{N_0}(\overline{\mathbb{R}^+})$ for a fixed integer $N_0 \geq 2$.

Then, the following holds, and can be proved by the same arguments of the proofs of Theorems 1.1-1.2.

Theorem 4.10. *Suppose that (\mathfrak{C}_1) is true. Then for each $N \in \{2, \dots, N_0\}$, all results of Theorems 1.1-1.2, except for (1.11)-(1.12) and (1.14), are true.*

5 An example

This section presents the decomposition (1.9) with a simple kernel. In this case, all coefficients h_l and p_l (with $l \in \mathbb{N}$) can be clearly given, from which we can easily see that for each $t \geq 0$ and each $l \geq 1$, $h_l(t) \neq 0$. So for each $N \geq 2$ and each $t \geq 0$, both heat-like terms and wave-like terms can be split away from the decomposition, and as N gets large, the principle part of the decomposition contains more and more terms.

To this end, we let

$$M(t) = \alpha e^{\lambda t}, \quad t \geq 0, \quad \text{with fixed } \alpha, \lambda > 0. \quad (5.1)$$

From (1.7)-(1.10), after direct computations, we find that when $N = 2$,

$$\begin{aligned} \Phi(t) &= \mathcal{P}_2(t) + \mathcal{W}_2(t) + \mathfrak{R}_2(t) \\ &= \left\{ e^{tA} + e^{tA} \left[-\alpha t A^{-1} + \left(\alpha - \alpha \lambda t + \frac{1}{2} \alpha^2 t^2 \right) A^{-2} \right] \right\} \\ &\quad - \alpha e^{\lambda t} A^{-2} + \mathcal{R}_2(t, -A)(-A)^{-3}, \quad t \geq 0, \end{aligned} \quad (5.2)$$

where

$$\mathcal{R}_N(t, \tau) = (-1)^N N! \int_0^t \tau e^{-\tau s} e^{\lambda(t-s)} \mathcal{F}_N(t, s) ds, \quad t, \tau \geq 0 \quad (5.3)$$

with

$$\mathcal{F}_N(t, s) := \sum_{\substack{\beta_1, \beta_2, \beta_3 \in \mathbb{N}, j \in \mathbb{N}^+ \\ \beta_1 + \beta_2 + \beta_3 = N, \\ \beta_1 \leq j, \beta_2 \leq j-1}} \frac{(-s)^{j-\beta_1} (t-s)^{j-1-\beta_2}}{(j-\beta_1)! (j-1-\beta_2)!} \lambda^{\beta_3} \alpha^j.$$

From (5.2), as well as (5.3), we see what follows: First, \mathcal{P}_2 and \mathcal{W}_2 shows clearly the heat-like effect and the wave-like effect respectively; Second, the remainder \mathfrak{R}_2 still has a complicated expression.

For the case that M is given by (5.1) and $N \geq 2$ is arbitrarily given, we can use (1.7)-(1.10) to get the decomposition of the flow with clearly expressed first two components, together with a complicated expressed remainder. Indeed, by (1.8) and (5.1), we have that for each $l \in \mathbb{N}$,

$$\begin{cases} h_l(t) = e^{\lambda t} (-1)^l \sum_{\substack{m, k \in \mathbb{N}, j \in \mathbb{N}^+ \\ k-m+2j=l+1, \\ m+k \leq l-1}} C_l^{l-j} \alpha^j \lambda^k \frac{t^m}{m!}, & t \geq 0, \\ p_l(t) = (-1)^{l+1} \sum_{\substack{m, k \in \mathbb{N}, j \in \mathbb{N}^+ \\ k-m+2j=l+1, \\ m+k \leq l+1}} C_l^{l-j+m} \alpha^j \lambda^k \frac{(-t)^m}{m!}, & t \geq 0. \end{cases}$$

Here, we use the conventional notation: $0^0 := 1$ and $\sum_{\emptyset} \cdot := 0$. From the above and (5.1), one can easily see that

$$h_l(t) \neq 0 \text{ for each } t \geq 0 \text{ and } l \geq 1. \quad (5.4)$$

The reason that remainders always have complicated expressions can be explained as follows: The remainders inherit the hybridity of heat and waves of the flow, no matter how big N is, and such hybridity cannot have a simple expression in terms of the operator A .

At last, we mention that such kind of initial data $\mathcal{P}_N(0)y_0$, under the action of the flow $\Phi(\cdot)$, may have three nontrivial components at positive time. More precisely, for each integer $N \geq 2$ and $T \in (0, 1)$, there is $j \in \mathbb{N}^+$ so that for each $t \in [T, 1/T]$, the following equality

$$\begin{aligned} \Phi(t)(\mathcal{P}_N(0)e_j) &= I_1 + I_2 + I_3 \\ &\triangleq \mathcal{P}_N(t)(\mathcal{P}_N(0)e_j) + \mathcal{W}_N(t)(\mathcal{P}_N(0)e_j) + \mathfrak{R}_N(t)(\mathcal{P}_N(0)e_j) \end{aligned}$$

has the following property

$$I_1, I_2, I_3 \neq 0. \quad (5.5)$$

(This is also true when $\mathcal{P}_N(0)e_j$ is replaced by $\mathcal{W}_N(0)e_j$ in the above statement.) In fact, it follows from (1.9) twice, as well as (1.10), that for each $t > 0$,

$$\begin{aligned}\mathfrak{R}_N(t) &= (\mathcal{P}_{N+1}(t) - \mathcal{P}_N(t)) + (\mathcal{W}_{N+1}(t) - \mathcal{W}_N(t)) + \mathfrak{R}_{N+1}(t) \\ &= p_N(t)e^{tA}(-A)^{-N-1} + h_N(t)(-A)^{-N-1} + \mathfrak{R}_{N+1}(t).\end{aligned}$$

Then by using (1.10) again, we see that for each $j \geq 1$,

$$\begin{cases} \mathcal{P}_N(t)(\mathcal{P}_N(0)e_j) &= e^{-t\eta_j} \left(1 + \sum_{l=0}^{N-1} p_l(t)\eta_j^{-l-1}\right) \left(1 + \sum_{l=0}^{N-1} p_l(0)\eta_j^{-l-1}\right) e_j, \\ \mathcal{W}_N(t)(\mathcal{P}_N(0)e_j) &= \left(\sum_{l=0}^{N-1} h_l(t)\eta_j^{-l-1}\right) \left(1 + \sum_{l=0}^{N-1} p_l(0)\eta_j^{-l-1}\right) e_j, \\ \mathfrak{R}_N(t)(\mathcal{P}_N(0)e_j) &= \left(h_N(t)\eta_j^{-N-1} + p_N(t)e^{-t\eta_j}\eta_j^{-N-1} + \mathcal{R}_{N+1}(t, \eta_j)\eta_j^{-N-2}\right) \\ &\quad \left(1 + \sum_{l=0}^{N-1} p_l(0)\eta_j^{-l-1}\right) e_j. \end{cases}$$

Since $\lim_{j \rightarrow +\infty} \eta_j = +\infty$, from the above and (5.4), (5.5) follows for some large j . (By a similar way, (5.5) can be verified in the case that $\mathcal{P}_N(0)e_j$ is replaced by $\mathcal{W}_N(0)e_j$.)

6 Conclusions and further comments

In this paper we have presented a decomposition for the flow generated by the equation (1.1), which reveals the hybrid parabolic-hyperbolic behavior of the flow. We have also described the nature of each of the components in the decomposition; that has been illustrated through an example.

A number of interesting issues could be considered in connection with the results and methods developed in this paper. Here, we briefly give some of them.

- *Smooth memory kernels.* It would be interesting to analyze whether (1.11), (1.12) and (1.14) hold under the assumption that $M \in C^\infty(\overline{\mathbb{R}^+}) \setminus \{0\}$.
- *Decomposition with infinite series.* It would be interesting to obtain a meaningful decomposition without the intervention of the third remainder term.
- *Space-dependent memory kernels.* The extension of the results of this paper to the space-dependent memory kernels $M = M(t, x)$ is open.
- *Other equations with memory.* It would be interesting to extend this decomposition to other models such as wave equations with memory kernels.

7 Appendix

Proposition 7.1. *For each $t \geq 0$ and $s \in \mathbb{R}$, it holds that*

$$\Phi(t)y_0 = y(t; y_0), \quad y_0 \in \mathcal{H}^s \tag{7.1}$$

and that $\Phi(t)$ belongs to $\mathcal{L}(\mathcal{H}^s)$.

Proof. Arbitrarily fix $s \in \mathbb{R}$. Since $\{e^{tA}\}_{t \geq 0}$ is a C_0 semigroup over \mathcal{H}^s , we can use a standard method (see for instance [18, Theorem 1.2 in Section 6.1, p. 184]) to show that for each $y_0 \in \mathcal{H}^s$, the solution $y(\cdot; y_0)$ belongs to the space $C(\overline{\mathbb{R}^+}; \mathcal{H}^s)$. This, along with (1.2), yields (7.1). Moreover, one can directly check that for each $t \geq 0$, $\Phi(t) \in \mathcal{L}(\mathcal{H}^s)$. This completes the proof. \square

The proof of Lemma 2.1 is given as follows.

Proof of Lemma 2.1. By standard density arguments, it suffices to show that for each $k \in \mathbb{N}$, the following property (\mathcal{E}_k) is true: For all kernels $\{M_l\}_{l=1}^j \subset C^\infty(\overline{\mathbb{R}^+})$,

$$\left| \frac{d^k}{dt^k} M_1 * \cdots * M_j(t) \right| \leq \mathcal{C}(j, k, t) \prod_{l \leq q(j, k)} \|M_l\|_{C^{p(j, k)}([0, t])} \prod_{l > q(j, k)} \|M_l\|_{C^{p(j, k)-1}([0, t])}, \quad t > 0. \quad (7.2)$$

Here and in what follows, we set $\prod_{\emptyset} \cdot := 1$ and let

$$\begin{cases} p(j, k) &:= \max\{k - j + 1, 1\}, \\ q(j, k) &:= \min\{k, j\}, \\ \mathcal{C}(j, k, t) &:= \sum_{l=j-1-k}^{j-1} \chi_{\mathbb{N}}(l) \frac{t^l}{l!}, \quad t > 0. \end{cases} \quad (7.3)$$

Now we will use the induction to prove the above (\mathcal{E}_k) for each $k \in \mathbb{N}$. To this end, we first check (\mathcal{E}_0) . Indeed, for each $j \in \mathbb{N}^+$ and each $\{M_l\}_{l=1}^j \subset C^\infty(\overline{\mathbb{R}^+})$, we have that

$$|M_1 * \cdots * M_j(t)| \leq \left(\prod_{l=1}^j \|M_l\|_{C([0, t])} \right) \int_0^t \int_0^{t_1} \cdots \int_0^{t_{j-1}} dt_j \cdots dt_1 = \frac{t^{j-1}}{(j-1)!} \prod_{l=1}^j \|M_l\|_{C([0, t])}.$$

This, along with (7.3), leads to (7.2) with $k = 0$. Therefore (\mathcal{E}_0) is true.

Next, we will show (\mathcal{E}_{k_0+1}) for any $k_0 \in \mathbb{N}$, under the assumption that (\mathcal{E}_k) holds for all $k \leq k_0$. For this purpose, we arbitrarily fix $k_0 \in \mathbb{N}$, $j \in \mathbb{N}^+$ and $\{M_l\}_{l=1}^j \subset C^\infty(\overline{\mathbb{R}^+})$. Since (7.2), with $j = 1$, holds clearly, we only need to focus on the situation that $j \geq 2$. There are only two possibilities for j : either $k_0 \leq j - 1$ or $k_0 \geq j$.

In the case when $k_0 \leq j - 1$, we have three observations: First, by direct computations, we find

$$\begin{aligned} \frac{d^{k_0+1}}{dt^{k_0+1}} M_1 * \cdots * M_j &= \frac{d^{k_0}}{dt^{k_0}} \left(\frac{d}{dt} M_1 * \cdots * M_j \right) \\ &= \frac{d^{k_0}}{dt^{k_0}} \left(M_1(0) M_2 * \cdots * M_j + M_1' * M_2 * \cdots * M_j \right) \\ &= M_1(0) \frac{d^{k_0}}{dt^{k_0}} M_2 * \cdots * M_j + \frac{d^{k_0}}{dt^{k_0}} M_2 * \cdots * M_j * M_1'. \end{aligned} \quad (7.4)$$

Second, we apply (\mathcal{E}_{k_0}) twice to find that for each $t > 0$,

$$\left| \frac{d^{k_0}}{dt^{k_0}} M_2 * \cdots * M_j(t) \right| \leq \mathcal{C}(j-1, k_0, t) \left(\prod_{2 \leq l \leq k_0+1} \|M_l\|_{C^1([0, t])} \right) \left(\prod_{l > k_0+1} \|M_l\|_{C([0, t])} \right); \quad (7.5)$$

$$\left| \frac{d^{k_0}}{dt^{k_0}} M_2 * \cdots * M_j * M'_1(t) \right| \leq \mathcal{C}(j, k_0, t) \left(\prod_{2 \leq l \leq k_0+1} \|M_l\|_{C^1([0,t])} \right) \left(\prod_{l > k_0+1} \|M_l\|_{C([0,t])} \right) \|M'_1\|_{C([0,t])}. \quad (7.6)$$

Third, from the third definition in (7.3), we see

$$\max \left\{ \mathcal{C}(j-1, k_0, t), \mathcal{C}(j, k_0, t) \right\} \leq \mathcal{C}(j, k_0+1, t), \quad t > 0. \quad (7.7)$$

Now, from (7.4), (7.5), (7.6) and (7.7), it follows that for each $t > 0$,

$$\begin{aligned} \left| \frac{d^{k_0+1}}{dt^{k_0+1}} M_1 * \cdots * M_j(t) \right| &\leq \mathcal{C}(j, k_0+1, t) \left(\|M_1\|_{C([0,t])} + \|M'_1\|_{C([0,t])} \right) \\ &\quad \times \left(\prod_{2 \leq l \leq k_0+1} \|M_l\|_{C^1([0,t])} \right) \left(\prod_{l > k_0+1} \|M_l\|_{C([0,t])} \right). \end{aligned}$$

This, along with (7.3), leads to (7.2) (with $k = k_0 + 1 \leq j$). Therefore, (\mathcal{E}_{k_0+1}) is true when $k_0 \leq j-1$.

In the case that $k_0 \geq j$, we see from (7.3) that

$$\begin{cases} p(j, k_0+1) = p(j-1, k_0) = p(j, k_0) + 1 = k_0 - j + 2; \\ q(j, k_0+1) = q(j, k_0) = q(j-1, k_0) + 1 = j. \end{cases}$$

Then by the similar arguments as those in (7.4)-(7.7), one can get that for each $t > 0$,

$$\begin{aligned} \left| \frac{d^{k_0+1}}{dt^{k_0+1}} M_1 * \cdots * M_j(t) \right| &\leq \mathcal{C}(j, k_0+1, t) \left(\|M_1\|_{C([0,t])} + \|M'_1\|_{C^{p(j,k_0)}([0,t])} \right) \\ &\quad \times \prod_{l=2}^j \|M_l\|_{C^{p(j-1,k_0)}([0,t])} \\ &= \mathcal{C}(j, k_0+1, t) \prod_{l=1}^j \|M_l\|_{C^{p(j,k_0+1)}([0,t])}. \end{aligned}$$

This, together with (7.3), leads to (7.2) (with $k = k_0 + 1 > j$). Therefore, (\mathcal{E}_{k_0+1}) holds in the case that $k_0 \geq j$.

Hence, we complete the proof of Lemma 2.1. \square

Acknowledgments. The authors would like to gratefully thank Dr. Huaiqiang Yu and Dr. Christophe Zhang for their valuable comments and suggestions.

The first and second authors were supported by the National Natural Science Foundation of China under grants 11971022 and 11801408, respectively.

The third author has been funded by the Alexander von Humboldt-Professorship program, the European Research Council (ERC) under the European Union's Horizon 2020 research and innovation programme (grant agreement No. 694126-DyCon), the grant MTM2017-92996 of MINECO (Spain), ELKARTEK project KK-2018/00083 ROAD2DC of the Basque Government, the Nonlocal PDEs: Analysis, Control and Beyond, AFOSR Grant FA9550-18-1-0242 and the Marie Skłodowska-Curie grant agreement No. 765579-ConFlex.

References

- [1] G. Amendola, M. Fabrizio and J. Golden. Thermodynamics of materials with memory: Theory and applications. Springer, New York, 2012.
- [2] L. Boltzmann. Zur theorie der elastischen nachwirkung. Wien. Ber. 70 (1874) 275–306.
- [3] L. Boltzmann. Zur theorie der elastischen nachwirkung. Wien. Ber. 5 (1878) 430–432.
- [4] C. Cattaneo. A form of heat conduction equation which eliminates the paradox of instantaneous propagation. Compute. Rendus. 247 (1958) 431–433.
- [5] F. Chaves-Silva, X. Zhang and E. Zuazua. Controllability of evolution equations with memory. SIAM J. Control Optim. 55 (2017) 2437–2459.
- [6] R. Christensen. Theory of viscoelasticity, an introduction. Academic Press, New York, 1982.
- [7] B. Coleman and M. Gurtin. Equipresence and constitutive equations for rigid heat conductors. Z. Angew. Math. Phys. 18 (1967) 199–208.
- [8] C. Dafermos. Asymptotic Stability in Viscoelasticity. Arch. Ration. Mech. Anal. 37 (1970) 297–308.
- [9] M. Fabrizio, C. Giorgi and V. Pata. A new approach to equations with memory. Arch. Ration. Mech. Anal. 198 (2010) 189–232.
- [10] X. Fu, J. Yong and X. Zhang. Controllability and observability of the heat equations with hyperbolic memory kernel. J. Differential Equations. 247 (2009) 2395–2439.
- [11] M. Gurtin and A. Pipkin. A general theory of heat conduction with finite wave speeds. Arch. Ration. Mech. Anal. 31 (1968) 113–126.
- [12] L. Hörmander. The Analysis of Linear Partial Differential Operators, Vol. 3, Springer-Verlag, 2007.
- [13] Q. Lü, X. Zhang and E. Zuazua. Null controllability for wave equations with memory. J. Math. Pures Appl. 108 (2017) 500–531.
- [14] J. Maxwell. On the dynamical theory of gases. Phil. Trans. Roy. Soc. London 157 (1867) 49–88.
- [15] M. Reed and B. Simon. Methods of Modern Mathematical Physics I: Functional Analysis, Academic Press, 1980.
- [16] M. Reed and B. Simon. Methods of Modern Mathematical Physics IV: Analysis of Operators, Academic Press, 1978.
- [17] L. Pandolfi. Linear systems with persistent memory: An overview of the bibliography on controllability. arXiv: 1804.01865v1 (2018), preprint.
- [18] A. Pazy. Semigroups of Linear Operators and Application to Partial Differential Equations. Springer-Verlag, New York, Inc., 1983.

- [19] V. Volterra. Sur les équations intégrales-différentielles et leurs applications. *Acta Math.* 35 (1912) 295–356.
- [20] V. Volterra. *Leçons sur les fonctions de lignes*. Gauthier-Villars, Paris, 1913.
- [21] G. Wang, Y. Zhang and E. Zuazua. Reachable subspaces, control regions and heat equations with memory. *arXiv:2101.10615v1*, preprint.